

Role of antioxidant therapy and cessation of smoking in treatment of infertile men.

Falah Mahdi Ali Al-khafaji* , Jawad kadham Al-janabi**

Mohammed Ali Ghadhban***

* Branch of Surgery, Urology field, FIBMS(Urology), Al-Qadissiya University - College of Medicine, Al-Diwaniya.** Public health laboratory, PhD- microbiology, department of health – Al-Diwaniya.***Department of Surgery, Urology field, FIBMS(Urology), Al-Diwaniya teaching hospital,Al-Diwaniya.

(Received 25/ 1 /2015 , Accepted 26 / 3 / 2015)

الخلاصة

الهدف من الدراسة : لتقييم دور علاج مضادات الأوكسدة وتوقف التدخين في مرضى عقم الرجال. المرضى وطرق إجراء الدراسة : أجريت الدراسة في شهر شباط 2012 إلى شهر ايلول 2013 . العدد الكلي للمرضى كان 152 مريض ، 35 مريض كانوا مصابين بعقم ثانوي، مدى مدة الزواج كان 1-8 سنوات، وكان مدى التدخين من 15-40 سيجارة في اليوم وعلى فترة من 2-15 سنة، وشملت الدراسة أنواع متعددة من العقم (قلة العدد، ضعف الحيامن، تشوه الحيامن، أو جميعهم). أما متلازمة فقد النطاف الكلي فكان مستبعد عن الدراسة. 90 مريض عولج بالمضادات بدون التوقف من التدخين (رفض ذاتي)، و 62 مريض عولج بالمضادات مع توقف التدخين. كل المرضى اجري لهم تحاليل الهرمونات في الدم.

النتائج : 27 مريض مصاب بضعف النطف أوقفوا التدخين ، وكان متوسط القدرة على الحركة 16,4 بالمئة وبعد العلاج أصبح 30,1 بالمئة ، 53 مريض مصاب بضعف النطف من الذين لم يوقفوا التدخين ، كان متوسط القدرة على الحركة 15,3 بالمئة وبعد العلاج أصبح 31 بالمئة . 23 مصاب بقلة عدد النطف أوقفوا التدخين كان متوسط العدد بالمليون 6,3 وبعد العلاج أصبح 15,2. 33 مريض مصابين بقلة عدد النطف لم يوقفوا التدخين كان متوسط العدد بالمليون 6,4 وبعد العلاج أصبح 9,2. أما بالنسبة لمرضى الحيامن المشوهة فكان 5 منهم قد أوقفوا التدخين وكان متوسط نسبة الشكل غير المشوه 1,8 بالمئة وبعد العلاج أصبح 12,4 بالمئة و 11 مريض مصابين بتشوهات الحيامن ممن لم يوقفوا التدخين كان نسبة النطف غير المشوهة 1,7 بالمئة وبعد العلاج أصبح 5,6 بالمئة.

الاستنتاج : استخدام مضادات الأوكسدة والتوقف عن التدخين في مرضى العقم ممن اجري لهم هورمونات الجنس وكانت طبيعية يعطي نتائج جيدة بالنسبة لعدد النطف وقدرتها على الحركة.

Abstract

Aim: To assess role of antioxidant therapy and cessation of smoking in infertile men .

Patients and Methods: This study was conducted at infertility unit at Al- Diwaniya gynecological hospital and private clinic from February 2012 to November 2013. Total number of patients was taken in study were 152. Thirty-five patients were secondarily infertile while 117 patients were primarily infertile. Range of duration of marriage was 1-8

years. Each man smoked from 15-40 cigarette per day for 2-15 years. The study was include patients presented with various seminal fluid analytic pictures (oligospermia, teratospermia, asthenospermia, OAT syndrome). Ninety patients were treated by antioxidants without cessation of smoking and 62 patients were treated by antioxidants with cessation of smoking.

Results: Regarding 27 asthenospermic ceased smoking patients (mean motility percentage before treatment is 16.4 and mean motility percentage after treatment is 30.1), 53 asthenospermic non ceased smoking patients (mean motility percentage before treatment is 15.3 and mean motility percentage after treatment is 31), 23 oligospermic ceased smoking patients (mean count per million before treatment is 6.3 and mean count per million after treatment is 15.2), 33 oligospermic non ceased smoking patients (mean count per million is 6.4 and mean count per million after treatment is 9.2). regarding 5 teratospermic ceased smoking patients (mean percentage of abnormal morphology before treatment is 1.8 and mean percentage of abnormal morphology after treatment is 12.4) and 11 teratospermic non ceased smoking patients(mean percentage of abnormal morphology before treatment is 1.7 and mean percentage of abnormal morphology after treatment is 5.6).

Conclusion: use of antioxidant and cessation of smoking in selected patients (normal serum hormones) gives good improvement regarding motility and count of sperms

Introduction

Infertility is inability of a couple to achieve conception or bring a pregnancy to term after one year or more of regular, unprotected sexual intercourse according to world health organization definition ⁽¹⁾.

It affected at least one in six couples ⁽²⁾. In about 50% of cases, male factors play a role in couples infertility ^(3,4).

The etiology of abnormal semen quality is poorly understood, and many physiological, anatomical, environmental, and genetic factors have been implicated ⁽⁵⁻⁸⁾.

The oxidative stress is mechanism resulted from these factors which is induced by reactive oxygen species(ROS) or free radicals, excessive ROS (superoxide anion, hydrogen peroxide, hydroxyl radical) can damage sperm by modifying cell functions or endanger cell survival or both ⁽⁹⁾.

ROS are products of normal cellular metabolism. During the enzymatic reduction of O₂ to produce energy, free radicals form as a byproduct ⁽¹⁰⁾.

The unique cellular structure of spermatozoa renders them particularly sensitive to oxidative stress in that the sperm cells lack significant cytoplasm and therefore contain only minimal amounts of

ROS scavenging enzymatic substances(catalase, superoxide dismutase(SOD), and glutathione peroxidase).

Sources of ROS :

1-leukospermia as a result of infection/ inflammation ⁽¹¹⁻¹⁹⁾.

2-exposure to industrial compounds(phthalate in plastic material used in food packaging), pesticides(contains lindane) ^(20,21).

3-cigarrete smoking ⁽²²⁻²⁶⁾.

4-alcohol intake ⁽²²⁻²⁶⁾.

5-high body temperature like fever ⁽²⁷⁾ or wearing tight fitting underwear⁽²⁸⁾.

Exposure to cigarette smoke generate high levels of ROS, directly increasing both seminal leukocyte concentration and seminal ROS generation ^(22,26).

Effects of smoking on testicular function: decreasing Semen quality and sperm fertilizing capacity and decreasing seminal levels of antioxidant enzymes ⁽²⁹⁾.

In addition to that the smoking increase production of norepinephrine which increases the conversion of testosterone to

estrogen causing decreased testosterone levels⁽³⁰⁾.

Nicotine-induced suppression of hypothalamic-pituitary-testicular axis that is one of the causes of smoking-induced adverse effects on male infertility through suppression of LH secretion⁽³¹⁾.

Antioxidants maintains a steady state of ROS in the seminal plasma(act as free radical scavengers). In addition, semen contain a variety of non-enzymatic antioxidant molecules such as vitamin C, vitamin E, glutathione, selenium, and carnitine⁽³²⁾. These antioxidants compensate for the loss of sperm cytoplasmic enzymes as the cytoplasm is extruded during spermiogenesis, which in turn, diminishes

endogenous repair mechanism and enzymatic defenses⁽³³⁾.

Patients and Methods

This prospective clinical trial was conducted at infertility unit at Al- Diwaniya gynecological hospital and private clinic from February 2012 to November 2013.

One hundred fifty two was the total number included in the study. Thirty-five patients were secondarily infertile while 117 patients were primarily infertile. Range of duration of infertility was 1-8 years. Each man smoked from 15-40 cigarette per day for 2-15 years.

The study was include patients presented with various seminal fluid analytic pictures (oligospermia, teratospermia, asthenospermia, OAT syndrome).

infertility duration / year	1-8
Number of cigarettes / day	15-40
Duration of smoking / year	2-15
Number of primary infertile men	117
Number of secondary infertile men	35

Azoospermia, varicocele and pyospermia were excluded from the study.

Table 1 : summarize the above data.

All their wives were consult their gynecologists.

Ninety patients were treated by antioxidants(mixture of zinc, selenium, L-Carnitine, L-arginine and Vitamin E) without cessation of smoking and 62 patients were treated by antioxidants with cessation of smoking.

Semen samples were collected by usual way (masterbation)and analyzed under microscopy. Serum Pituitary- Testicular hormones : LH, FSH, Testosterone were measured.

Results

The table below reveals results of the study after prescribing antioxidant treatment with and without cessation of smoking. regarding 27 asthenospermic ceased smoking patients (mean motility percentage[grade A] before

treatment is 16.4 and mean motility percentage[grade A] after treatment is 36.1), 53 asthenospermic non ceased smoking patients(mean motility percentage[grade A] before treatment is 15.3 and mean motility percentage [grade A] after treatment is 31), 23 oligospermic ceased smoking patients(mean count per million before treatment is 6.3 and mean count per million after treatment is 15.2), 33 oligospermic non ceased smoking patients (mean count per million is 6.4 and mean count per million after treatment is 9.2). regarding 5 teratospermic ceased smoking patients (mean percentage of normal morphology before treatment is 1.8 and mean percentage of normal morphology after treatment is 12.4)and 11 teratospermic non ceased smoking patients(mean percentage of normal morphology before

treatment is 1.7 and mean percentage of normal morphology after treatment is 5.6).

	Mean	N	Std. Deviation	P value
Asthenopretreatment ceased	16.4074	27	9.26224	<0.001
asthenoposttreatment ceased	36.1852	27	11.61245	
asthenopretreatment n ceased	15.3396	53	10.17906	<0.001
asthenopostnonceased	31.0943	53	11.82121	
oligopreceased	6.3348	23	3.57995	<0.001
oligopostceased	15.2609	23	7.57704	
oligoprenonceased	6.4576	33	4.04529	0.001
oligopostnonceased	9.2636	33	5.07068	
nonteratopreceased	1.8000	5	.83666	0.030
nonteratopostceased	12.4000	5	6.58027	
nonteratoprenonceased	1.7273	11	.78625	0.076
nonteratopostnonceased	5.6364	11	6.56160	

Table 2: results after treatment by antioxidant with or without cessation of smoking.




Figure1: improvement of motility(grade A) in patients stop smoking




Figure2: improvement of motility(grade A) in patients not stop smoking




Figure3: improvement of sperm count in patients stop smoking




Figure4: improvement of sperm count in patients not stop smoking




Figure5: improvement of sperm morphology in patients stop smoking




Figure6: improvement of sperm morphology in patient not stop smokin

Discussion

As we know, infertility is inability to get pregnancy after one year or more of regular sexual intercourse without contraceptive procedures and this is may be due to a lot of reasons (genetic , environmental , physiological causes) or may be idiopathic⁽³⁴⁻³⁷⁾.

From the environmental factors is oxidative stress and smoking as they mentioned in many studies like (Agarwal et al , Saleh et al, Chen H et al)^(23, 38,39).

Oxidative stress is due to increased level of reactive oxygen species or antioxidant disability within semen⁽⁴⁰⁾.

Sperms are susceptible to oxidative stress for 2 reasons, firstly high polyunsaturated fatty acids and secondly deficiency of intracellular antioxidant enzymatic protection , and this is lead to DNA damage.

Normally epididymis contain enzymatic and non-enzymatic antioxidants protect sperms, enzymatic like dismutase, catalase, non-enzymatic like vitamin C , E , and many.

A lot of studies have carried out to establish beneficial effects of antioxidant in improving sperm quality and quantity.

From these studies: Fraga et al were used Vitamin C, Kodama et al were used Glutathione, Keskes-Amma et al were used Selenium, Costa et al were used L-carnitine, Kynaston et al were used Zinc, Giovenco et al were used Vitamin E . Semen of smoker patient contain high proportion of malformed sperm⁽⁴¹⁻⁴³⁾.

By-product of nicotine present in the semen of smokers have been found to reduce motility of sperm and their fertilization capacity⁽⁴³⁾.

In our study, we found that abnormal sperm count and motility in infertile smoker whether ceased or not were improved after treatment with antioxidant and this is due to the opposite effect on oxidative stress on sperm count and motility where p value is significant in asthenospermic and oligospermic patients after treatment especially more in patients where ceased smoking.

Most of urologists think that presence of reactive oxygen species is existing normally in every man but because of imbalance between them and scavengers , the reactive oxygen species will offend and cause jeopardy to sperms^(38,44).

In our study, were we use antioxidants inform of Vitamin E (α -tocopherol), Vitamin C, selenium , L-carnitine , to every patient and as we know effects of these antioxidants (suppress lipid peroxidation in testicular mitochondria and reverse the harmful effect of oxidative stress and then maintain spermatogenesis and maturation of the sperms.

In our study, were we instruct all patients to cease smoking but from 152 patients only 55 patients cease and those were got better improvement due to dangerous effect of smoking because smoking increase oxidative stress and cause more lipid peroxidation and inhibition of DNA synthesis
(34,39,45-47)

In teratospermic patients , were we found that p value is insignificant and this may be due to little number of patients and this lead to more standard errors.

We recommend from our study : to use of antioxidant and cessation of smoking in selected patients (normal serum hormones) may get good improvement regarding motility and count of sperms and also recommend selection of large number of teratospermic patients to get better results in future studies.

Conclusion

use of antioxidant and cessation of smoking in selected patients (normal serum hormones) gives good improvement regarding motility and count of sperms

References

1. Rowe PJ, Comhaire FH, Hargreave TB, Mahmoud AMA. WHO Manual for the Standardized Investigation, Diagnosis and Management of the Infertile Male. Cambridge University Press, Cambridge, 2000.
2. Taylor A. ABC of subfertility: extent of the problem. *BMJ* 2003; 327: 434–436
3. Singh K, Jaiswal D. Human male infertility: a complex multifactorial phenotype. *Reprod Sci.* 2011; 18(5): 418-425.

4. Farhi J, Ben-Haroush A. Distribution of causes of infertility in patients attending primary fertility clinics in Israel. *Isr Med Assoc J.* 2011; 13(1):51-54.
5. Sharlip ID, Jarow JP, Belker AM et al. Best practice policies for male infertility. *Fertil. Steril.* 2002; 77: 873–82.
6. Vine MF. Smoking and male reproduction: A review. *Int. J. Androl.* 1996; 19: 323–37.
7. Auger J, Eustache F, Andersen AG et al. Sperm morphological defects related to environment, lifestyle and medical history of 1001 male partners of pregnant women from four European cities. *Hum. Reprod.* 2001; 16: 2710–7.
8. Kenkel S, Rolf C, Nieschlag E. Occupational risks for male fertility: An analysis of patients attending a tertiary referral centre. *Int. J. Androl.* 2001; 24: 318–26.
9. Agarwal A, Saleh R A, “Role of oxidants in male infertility: rationale, significance, and treatment”, *Urol. Clin. North Am.* (2002);29,(4): pp. 817–827.
10. Tremellen K. Oxidative stress and male infertility: A clinical perspective. *Hum. Reprod. Update* 2008; 14: 243–58.
11. Camejo MI, Segnini A, Proverbio F. Interleukin-6 in seminal plasma of infertile men, and lipid peroxidation of their sperm. *Arch. Androl.* 2001; 47: 97–101.
12. Martinez P, Proverbio F, Camejo MI. Sperm lipid peroxidation and pro-inflammatory cytokines. *Asian J. Androl.* 2007; 9: 102–7.
13. Nandipati KC, Pasqualotto FF, Thomas AJ, Agarwal A. Relationship of interleukin-6 with semen characteristics and oxidative stress in vasectomy reversal patients. *Andrologia* 2005; 37: 131–4.
14. Rajasekaran M, Hellstrom WJ, Naz RK, Sikka SC. Oxidative stress and interleukins in seminal plasma during leukocytospermia. *Fertil. Steril.* 1995; 64: 166–71.
15. Sanocka D, Jedzejczak P, Szumala-Kaekol A, Fraczek M, Kurpisz M. Male genital tract inflammation: The role of selected interleukins in regulation of pro-oxidant and antioxidant enzymatic substances in seminal plasma. *J. Androl.* 2003; 24: 448–55.
16. Baker MA, Krutskikh A, Aitken RJ. Biochemical entities involved in reactive oxygen species generation by human spermatozoa. *Protoplasma* 2003; 221: 145–51.
17. Fisher H, Aitken R. Comparative analysis of the ability of precursor

- germ cells and epididymal spermatozoa to generate reactive oxygen metabolites. *J. Exp. Zool.* 1997; 277: 390–400.
18. Latchoumycandane C, Mathur PP. Induction of oxidative stress in the rat testis after short-term exposure to the organochlorine pesticide methoxychlor. *Arch. Toxicol.* 2002; 76: 692–8.
19. Latchoumydecane C, Chitra KC, Mathur PP. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) induces oxidative stress in the epididymis and epididymal sperm of adult rats. *Arch. Toxicol.* 2003; 77: 280–4.
20. Agarwal DK, Maronpot RR, Lamb JC IV, Kluwe WM. Adverse effects of butyl benzyl phthalate on the reproductive and hematopoietic systems of male rats. *Toxicology* 1985; 35: 189–206.
21. Kasahara E, Sato EF, Miyoshi M et al. Role of oxidative stress in germ cell apoptosis induced by di(2-ethylhexyl)phthalate. *Biochem. J.* 2002; 365: 849–56.
22. Arabi M, Mosthagi H. Influence of cigarette smoking on spermatozoa via seminal plasma. *Andrologia* 2005; 37: 119–24.
23. Saleh RA, Agarwal A, Sharma RK, Nelson DR, Thomas AJ. Effect of cigarette smoking on levels of seminal oxidative stress in infertile men: A prospective study. *Fertil. Steril.* 2002; 78: 491–9.
24. Vine MF. Smoking and male reproduction: A review. *Int. J. Androl.* 1996; 19: 323–37.
25. Mostafa T, Tawadrous G, Roaia MM, Amer MK, Kader RA, Aziz A. Effect of smoking on seminal plasma ascorbic acid in infertile and fertile males. *Andrologia* 2006; 38: 221–4.
26. Sofikitis N, Miyagawa I, Dimitriadis D, Zavos P, Sikka S, Hellstrom W. Effects of smoking on testicular function: Semen quality and sperm fertilizing capacity. *J. Urol.* 1995; 154: 1030–4.
27. Carlsen E, Andersson AM, Petersen JH, Skakkebaek NE. History of febrile illness and variation in semen quality. *Hum. Reprod.* 2003; 18: 2089–92.
28. Evers CH, Tiemessen JL, Bots RS. Tight-fitting underwear and sperm quality. *Lancet.* 1996; 347: 1844–5.
29. Pasqualotto FF, Umezu FM, Salvador M, Borges E, Sobreiro BP, Pasqualotto EB.
30. Pasqualotto F F, Lucon A M, Sobreiro A M, Pasqualotto E B, Arap S, “Effects of medical therapy, alcohol, smoking, and endocrine disruptors on male infertility”, *Rev. Hosp. Clin. Fac. Med. Sao Paulo* (2004);59;(6): pp. 375–382.
31. Funabashi T, Sona A, Mitsushima d and Kimura F. Nicotine Inhibit Pulsatile LH secretion in human males but not in human females and tolerance to this nicotine effect loss within one week of quitting smoking. *J Clin Endocrinol Metab.* 2005; 90: 3908- 3913.
32. Agarwal A, Nallella K P, Allamaneni S S, Said T M, “Role of antioxidants in treatment of male infertility: an overview of the literature”, *Reprod. Biomed. Online* (2004);8;(6): pp. 616–627.
33. Aitken R J, Clarkson J S, Fishel S, “Generation of reactive oxygen species, lipid peroxidation, and human sperm function”, *Biol. Reprod.* (1989);41;(1): pp. 183–197.
34. Makker K, Agarwal A, Shaerma R: Oxidative stress & male infertility. *Indian J*, 2009; 129:357-367.
35. Aitken RJ, Baker MA, De Iuliis GN, Nixon B: New insights into sperm physiology and pathology. *Handb Exp Pharmacol*, 2010; 198:99-115.
36. Aitken RJ, Baker MA, De Iuliis GN, Nixon B: New insights into sperm physiology and pathology. *Handb Exp Pharmacol*, 2010; 198:99-115.
37. 20. Sikka SC: Oxidative stress and role of antioxidants in normal and abnormal sperm function. *Front Biosc J*, 1996; 1:78–86.
38. Agarwal A, Makker K, Sharma R: Clinical relevance of oxidative stress in male factor infertility: an update. *Am J Reprod Immun*, 2008; 59:2–11.
39. Chen H, Liu J, Luo L, Baig MU, Kim JM, Zirkin BR: Vitamin E, aging and Leydig cell steroidogenesis. *Exp Gerontol*, 2005; 40: 728–736.
40. Fujii J, Iuchi Y, Matsuki S, Ishii T. Cooperative function of antioxidant and redox systems against oxidative stress in male reproductive tissues. *Asian J Androl* 2003;5:231–242..
41. Ramlau-Hansen CH et al. Is smoking a risk factor for decreased semen quality? A cross-sectional analysis. *Hum Reprod.* 2007 Jan;22(1):188-96
42. Gaur DS, Talekar M, Pathak VP. Effect of cigarette smoking on semen quality of infertile men. *Singapore Med J.* 2007 Feb;48(2):119-23.
43. Schilling K et al. Prevalence of behaviour-related fertility disorders in a clinical sample: results of a pilot study. *Arch Gynecol Obstet.* 2012 Nov;286(5):1307-14. doi: 10.1007/s00404-012-2436-x.
44. Saalu LC: The incriminating role of reactive oxygen species in idiopathic male infertility: an evidence based evaluation. *Pak J Biol Sci*, 2010; 13: 413-422.

45. Mancini A, Festa R, Silvesterini A, Nicolotti N, Di Donna V, La Torri G, Pontecorvi A, Meucci E: Hormonal regulation of total antioxidant capacity in seminal plasma. *J Androl*, 2009; 30:534-540.
46. 25. Agarwal A, Saleh RA, Bedaiwy MA: Role of reactive oxygen species in the pathophysiology of human reproduction. *Fertil Steril*, 2003; 79:829–843.
47. Desai N, Sharma R, Maker K, Sabnegh E, Agarwal A: Physiological and pathological levels of reactive oxygen species in neat semen of infertile men. *Fertil Steril*, 2009; 92:1626–1631.