



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics



Biological Sciences

- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics
- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

**Indexed in Elsevier Bibliographic Database
(Scopus and EMBASE)**

SCImago Journal Rank 0.288

Impact factor 2.958*

Chemical Abstracts
Service (www.cas.org)



A division of the American Chemical Society

CODEN IJPBJ2



Elsevier Bibliographic databases (Scopus & Embase)

SNIP value – 0.77

SJR - 0.288

IPP - 0.479


SNIP – Source normalised impact per paper

SJR – SCImago Journal rank

IPP – Impact per publication

Source – www.journalmetrics.com

(Powered by scopus (ELSEVIER))



LUND
UNIVERSITY



JACKSONVILLE STATE UNIVERSITY

Jacksonville State University
Houston Cole Library
USA (Alabama)



UNIVERSITY OF
OXFORD

Oxford, United Kingdom

INDEX COPERNICUS
INTERNATIONAL

*And indexed/catalogued in
many more university*



*Instruction to Authors visit www.ijpbs.net

For any Queries, visit "contact" of www.ijpbs.net



EFFECT OF IRAQI ETHANOLIC PROPOLIS EXTRACTS ON THE REACTIVE NITROGEN SPECIES AND GENOTYPE ALTERATION IN DIABETIC NEPHROPATHY RATS

AMNA N. ZGHAIR¹ AND FERDOUS A. JABIR^{*,2}

¹Chemistry Dept., College of Education, Al-Qadisiyah University, Iraq

²Biochemistry Dept., College of Medicine, Al-Qadisiyah University,, Iraq

ABSTRACT

The purpose of this research is to identify the effect of the Iraqi extract ethanolic propolis (EEP) on the nitrogen radicals and most important of which Peroxynitrite and nitric oxide, in rat laboratory infected kidney failure as a result of diabetes induce by streptozotocin (60 mg / kg), as well as its effect on the candidate genes in diabetic nephropathy are glucose transporter1 (GLUT1) and nephrin promoter were found local EEP cannot reduced genotoxicity for STZ, but an effective influence on the free radicals where found a decrease in the concentration of Peroxynitrite radical was detected by detection of Nitrotyrosine (3- NT) by (ELISA), where the concentration increasing in diabetic groups while groups that treated by local EEP showed a decrease in the concentration of (3-NT) compared with control group, which indicates the low concentration of Peroxynitrite. As well as the results show the effect of local EEP on the low concentration of nitric oxide, and reduce the level of glucose in the blood, kidney function (creatinine, uric acid) and the presence of an improvement in the pathological changes of kidney tissues in the treated groups compared to the diabetic, this indicates that propolis is a strong immunological factors and antioxidant. The results we have obtained through this study to conclude that propolis cannot reduce the genotoxicity induced STZ but has immunological properties through its effect on reducing the concentration of 3-NT and NO[•] in Pre-treated and Post-treated groups and this is due to contain rich in flavonoids, phenolic acid, and others. It also has anti-oxidant properties through its effect anti-hyperglycemia and his effect moderate on kidney function by reducing the concentration of creatinine and uric acid in the blood through the improvement of histological changes in the kidney of rats with diabetes induced by STZ.

KEYWORD: GLUT1, Nephrin promoter, EEP, 3-NT, Peroxynitrite and Mutation.



FERDOUS A. JABIR

Biochemistry Dept., College of Medicine, Al-Qadisiyah University,, Iraq

ferdousabbas@gmail.com

*Corresponding author

INTRODUCTION

Diabetes is characterized by the absence or relative lack of insulin and this has the effect in increasing the level of glucose in the blood, it is classified to insulin-dependent diabetes mellitus (IDDM); called Type I, non-insulin-dependent diabetes mellitus (NIDDM); called Type II and other specific types associated with genetic defects in beta cell function⁽¹⁾. Diabetic nephropathy is a clinical syndrome characterized by the decline in glomerular filtration rate and excessive deposition of extracellular matrix proteins^(2,3). Oxidizing free radicals consist of two classes are reactive oxygen species (ROS) and reactive nitrogen species (RNS). Radicals of nitrogen common is nitric oxide (NO \cdot) and Peroxynitrite (ONOO \cdot)⁽⁴⁾. In vivo, peroxynitrite consist in macrophages, and endothelial cells, platelets, white blood cells, nerve cells, and the other by the reaction between superoxide (\cdot O₂ \cdot) and the radical of the nitric oxide (NO \cdot). Peroxynitrite is a species of short life (about 10 ms in vivo)⁽⁵⁾. Attributed of cytotoxicity to the nitric oxide NO \cdot and peroxynitrite ONOO \cdot radicals. Peroxynitrite reacts with the lipids, DNA and proteins cause oxidative damage and other series of reactions caused by free radicals. Peroxynitrite can cause DNA damage through oxidative modifications in each of the nucleobases and the backbone sugar-phosphate^(6,7). Nitrogen free radicals play a role in diabetic nephropathy where there is an accumulation of evidence supporting the key role of each of NO, superoxide (\cdot O₂ \cdot) and peroxynitrite, in the pathogenesis of diabetes and diabetic complications^(8,9,10). Several studies have been proposed that increased oxidative stress, and the paths influential formation of peroxynitrite such as poly-ADP ribosyl polymerase (PARP) is involved in causing the disease diabetes microvascular nephropathy^(11,12). For the diabetic nephropathy effect on the genes have been proposed, where many of the metabolic pathways and related categories of genes as candidates to play a role in genetic susceptibility to diabetic nephropathy. Genes that give susceptibility to diabetic nephropathy can be soliciting in different ways. It is candidate genes in diabetic

nephropathy: Glucose transporter 1 gene (GLUT1): mainly facilitates the transfer of glucose in mesangial cells⁽¹³⁾. The experiments indicate that GLUT1 may be associated with impairment of renal glomeruli (glomerulopathy)⁽¹⁴⁾. Increase GLUT1 occur relatively quickly after the onset of diabetes, in addition, high levels of sugar in stimulating insulin growth factor1 (IGF-1) glucose uptake in cells mediated mesangial⁽¹⁵⁾, indicating that glucose can stimulate activity glucose transporters individual, as well as gene expression of GLUT1 in these cells. Where the expression of GLUT1 gene leads to the morphology features of diabetic nephropathy and the emergence of this gene GLUT1 be related to the development of diabetic nephropathy. Nephron promoter gene: Nephron is a protein a recently identified, which is a key element in the slit diaphragm, which has a major role in the nomination of the plasma. Mutations in the genes of the human nephron are the result in the absence of slit diaphragm, massive proteinuria, and killer disorder known as congenital nephrotic syndrome of the Finnish type⁽¹⁶⁾. Decreases in nephron gene expression in diabetic nephropathy refer to the organization on the transcriptional level will weaken. The antioxidants have an important role against free radical damage, and extremely important to maintain optimum health. Where free radicals cause damage to cells and therefore have a key role in many diseases⁽¹⁷⁾. Fortunately, free radicals are controlled naturally by configuring the useful compounds known as antioxidants different. Propolis is a strong antioxidant, scavenges free radicals directly⁽¹⁸⁾, which contains the components of a typical phenolic flavonoids, phenolic acids and esters⁽¹⁹⁾. EEP led to lower levels of blood glucose, fructose amin, Malondialdehyde (MDA), and nitric oxide (NO), total cholesterol (TC), triglyceride (TG), low-density lipoprotein cholesterol, low density lipoprotein cholesterol in serum of fasting mice. And increase the level of serum cholesterol, high-density lipoprotein cholesterol (HDL-C)⁽²⁰⁾. Streptozotocin (STZ) 2-deoxy -2 - ([methyl (nitrous oxide) amino] carbonyl amino) - β -D- glucopyranose. (STZ; -

N nitro derivative of glucosamine) induce experimental diabetes in rats and mice⁽²¹⁾. Which lead to the degeneration of the islets of Langerhans in the beta cells⁽²²⁾.

MATERIALS AND METHODS

Was used 60 rats Sprague-Dawley weighing between 150 ± 10 g and housing in the house of the animal in the College of Veterinary Medicine at the University of Al-Qadisiyah were divided animals into four groups (control), (diabetes), (pre-treated) and (post-treated) groups. The induction of diabetes in rats after fasting overnight before intraperitoneal injection by a single dose by (STZ, Sigma-Aldrich) at a dose of (60 mg / kg of body weight). Followed by high blood sugar in the rats over a period of up to 72 hours, using test strips glucose⁽²³⁾, where the level of glucose in the blood of more than 250 mg /dl guide on diabetes⁽²⁴⁾. The experiment continued for 60 days. In the last day were sacrificed animals and collecting serum samples and kidney tissue samples for the purpose of testing them.

Preparation of local ethanolic extract propolis

Local EEP Was prepared extract ethanolic by way Krell⁽²⁵⁾. Determination of mutations of GLUT1 and nephrin promoter genes in kidney tissue by polymerase chain reaction -single strand confirmation polymorphism PCR-SSCP technique^(26,27). Primers have been designed by National Center for Biotechnology Information(NCBI) online: primers of GLUT1 gene were: '5GACTTGGAGGAAGCAGCAAAG3' (forward) and '5ATCTTCGGGTTTGGACCCAG 3' (Reverse) the product length size is 691 bp.

Primers of nephrin promoter gene were: 5'GGGCCTAGACAAGTGTTGCT3'(forward) and GTGTTCCCGGTAGAGCGAAA(Reverse) the length size is 387bp. Determination of the concentration serum 3-nitro tyrosine (3- NT) in the rats by using technology immunohistochemistry (ELISA) by method quantitative sandwich enzyme⁽²⁸⁾. Determination of serum(NO),uric acid and creatinine in rats by spectrophotometric method^(29,30, 31).

Histopathological examination

kidney tissues preserved in 10% neutral formalin buffer solution, after fixation, the tissue was trimmed and the specimen were washed with saline for (1-2hrs) and transferred to following Steps:1. Dehydration: Specimen were passed through ascending grades of ethanol alcohol (70%, 80%, 90%, 100%). For 1 hour in each concentration, 2. Clearing: Two solutions of xylol commonly used for clearing. The specimens rested 1 hour in each step, 3. Impregnation with Paraffin wax, 4. Blocking. 5. Sectioning, 6. Staining with hematoxilin & eosin.^(32,33,34). All groups read the data statistically by the program SPSS, version 17 software (2010). Test methods include a way ANOVA one-way and two-way to make comparisons between groups below the level of probability ($p < 0.05$). Mean \pm standard error is used to express all data.

RESULTS AND DISCUSSION

The figures (3) and (4) show the results that we have obtained by PCR-SSCP technique a genetic mutation in certain locations of genes GLUT1 and Nephrin promoter respectively.




Figure 1

Shows genetic mutation in the GLUT1 gene in diabetes group induced by STZ

where M refers to the parameter 100bp DNA Ladder, either numbers (1-3) samples G1(control), (4-6) G2 samples (Diabetic), (7-9) G3 samples (Pre-treated), (10-12) G4 samples (Post-treated).




Figure 2

Shows genetic mutation of the Neph rin promoter gene shows in diabetes groups induced by STZ

where M refers to the parameter 100bp DNA Ladder, either numbers (1-3) samples G1 (control group), (4-6) samples G2 (Diabetic), (7-9) G3 samples (Pre-treated), (10-12) G4 samples (Post-treated).

Table1
Number of lanes according to genotype of GLUT1 and Nephtrin promoter genes.

Genes name	Samples	Genotype	No. lanes
GLUT1	(1,2,8)	AA	4
	(5,6,7,11)	AB	5
	(4,9,10,12)	AC	0
Nephtrin promoter	(1,3,4,5,9,11,7)	AA	3
	(2)	AB	4
	(6,8,10)	AC	4
	(12)	AD	2

The results indicated of a genetic mutation in certain locations of genes GLUT1 and Nephtrin promoter in infected groups with diabetes induced by STZ in each of G2, and G3 and G4. The table(2) shows the difference in the four groups in the number of lanes this difference indicates that for genetic mutation, suggesting a lack of effect of a substance on the EEP to reduce material STZ genotoxicity.

Cytotoxic effects of STZ rely on DNA alkylation by the work of a particular site with DNA bases⁽³⁵⁾ and by free radicals generated during metabolism STZ⁽³⁶⁾. Therefore local EEP not effect on the mutation that induced STZ but local EEP decreases from peroxy nitrite and nitric oxide radicals. the table (2) and figure (3) show the changes in the body weight after diabetes stimulation.

Table2
The effect of local EEP on the body weight in diabetic nephropathy male rats.

Day	Group1 (control) Means \pm SE	Group2 (Diabetes) Means \pm SE	Group3 (pre-treatment) Means \pm SE	Group3 (post-treatment) Means \pm SE
1	156.9 \pm 1.44	157.5 \pm 1.24	156.45 \pm 1.7	156.3 \pm 1.53
6	168 \pm 1.87	169.88 \pm 2.09	167.8 \pm 1.83	168 \pm 2.97
12	186.7 \pm 2.39	184.22 \pm 1.82	184.22 \pm 1.82	180.9 \pm 1.9
18	201.3 \pm 1.3	202.6 \pm 1.36	203.3 \pm 1.96	195.4 \pm 3.94
24	218.8 \pm 3.5	218.3 \pm 3.33	219.5 \pm 4.4	215.8 \pm 2.9
30	239.1 \pm 1.4	220.8 \pm 4.02	225.5 \pm 4.24	230 \pm 3.4
36	252.8 \pm 2.37	229.5 \pm 1.84	234.2 \pm 1.89	239.3 \pm 2.06
42	276 \pm 1.24	235 \pm 1.94	242 \pm 2.05	252 \pm 2.57
48	307.3 \pm 3.96	243.15 \pm 1.57	256.1 \pm 5.15	288.5 \pm 3.08
54	339.7 \pm 1.71	258.1 \pm 1.3	270.5 \pm 2.27	319.5 \pm 3.33
60	355.5 \pm 2.66	255.8 \pm 2.1	295 \pm 2.2	329.8 \pm 4.04




Figure 3
The effect of EEP on the increase body weight in diabetic nephropathy induced by STZ in male rats.

After diabetes stimulation began to show changes in body weight, where the G1 in the case of a continuous increase gradually, while the G2 be in the case of a decrease in body weight significantly compared to the G2. The G3 and G4 show a decrease in body weight compared to the G1 but after treatment by local EEP showed a significant improvement

in body weight as in the table (2), propolis has a strong anti-oxidant and the scavenging effect against free radicals⁽³⁷⁾, this result suggests that propolis may improve metabolism troubled associated with diabetes. Either the table (3) and figure (4) show the effect of local EEP on the level of blood glucose in diabetic nephropathy rats.

Table 3
The effect of local EEP on the level of glucose in diabetic nephropathy male rats.

Day	Group1 (control)	Group2 (Diabetes)	Group3 (pre-treatment)	Group3 (post-treatment)
24	89.1±1.13	87.7±2.13	87.8±2.23	92.2±2.3
30	90.7±1.45	534±16.34	288.4±15.17	371±28.08
36	89.1±1.88	511.7±18	348.7±25.91	305.6±22.93
42	92.2±2.3	541.7±13.24	311.6±19.97	288.5±16.03
48	87.7±2.13	539.6±15.37	488.4±8.25	282.4±10.61
54	91±1.11	520.1±15.9	296.8±8.87	255.4±11.97
60	92.1±1.92	528.4±11.33	315.81±13.94	267.9±15.66




Figure 4
Effect of local EEP on the level of blood glucose in the diabetic nephropathy male rats.

Where there was a significant increase in blood glucose levels ($P < 0.05$). It is estimated that the effect of EEP where the reduction in glucose in the blood clearly in G3 and G4 comparison with the (G2). On the other hand, the table (3), explains the low level of blood glucose in the G4 rats more than G3 rats, but were significantly higher than the control group ($P < 0.05$). These results are consistent

with⁽³⁸⁾, which showed that the local EEP in rats had an impact hypoglycemic effectively. The presence of flavonoids and polyphenolic components of active ingredients effective anti-oxidant, strengthen systems antioxidant defense in the tissues of the pancreas⁽³⁹⁾. The figure(5) show effect of local EEP on the concentration of (3- NT).




Figure 5

The effect local EEP on the concentration of (3-NT) in diabetic nephropathy male rats.

where 3-NT concentration increased in diabetic nephropathy rats (G2) ($P < 0.05$) compared with G1, so indicates increased in oxidative stress and peroxynitrite generation. As the effect of local EEP it has been shown that it leads to a reduction in the concentration of 3-NT in G4, while G3 a low concentration of

3-NT, but more concentration than the G4, but the decline in these groups are not up to the G1 as shown in table (4). These results are consistent with^(40,41). Either figure(6) explain the effect of local EEP on the concentration of serum nitric oxide (NO).




Figure 6

The effect of local EEP on the concentration of NO in diabetic nephropathy male rats.

It has been found that there is a significant increase in the concentration of NO in a diabetic rats (G2), compared with the G1 ($P < 0.05$), while the G3 and G4 observed a decrease in the level of NO compared with a diabetic nephropathy rats, but still

significantly higher than the control group ($P < 0.05$), as shown table (4). These findings are consistent with⁽³⁸⁾. The effect of EEP on the level of uric acid in the rats serum show in figure (7).




Figure 7

The effect of Iraqi EEP on the level of uric acid in the diabetic nephropathy male rats.

During this study found a significant increase in the level of uric acid in the serum of diabetic rats (G2) compared with control rats (G1), ($P < 0.05$) but G3 and G4, there is an improvement in the level of uric acid in the blood, where it is significantly reduced

compared to G2 ($P < 0.05$), but much more than the G1 ($P < 0.05$), as shown in table (4). These findings are consistent with ⁽³⁶⁾. And the figure (8) explains the effect of local EEP on the level of creatinine in the serum rats.




Figure 8

The effect of EEP on the level of creatinine in the serum of diabetic nephropathy male rats.

there was a significant increase in the levels of serum creatinine in diabetic rats G2 compared to G1 ($P < 0.05$), but the groups treated by local EEP G3 and G4 respectively, it was observed that the local EEP works on a decrease in the level of creatinine ($P < 0.05$), this result is consistent with the

research^(38,42), it was noted that an increase in the level of creatinine in diabetic nephropathy rats (G2) refers to the development of diabetic nephropathy as show in table (3).

Table 3
The effect of EEP on the biochemical measurements in the serum of male rats.

Groups	Conc. (3-NT) ng/ml	Conc. NO μ mole/l	Conc. Uric acid mg/dl	Conc. Creatinine mg/l
Group1 control	1.33 \pm 0.499	6.04 \pm 0.31	2.7 \pm 0.44	0.8 \pm 0.62
Group2 diabetic	20.28 \pm 0.436	22.32 \pm 0.66	9.1 \pm 0.25	3.85 \pm 0.76
Group3 pre-treated	11.45 \pm 1.46	10.78 \pm 0.15	5.7 \pm 0.39	2.24 \pm 0.29
Group4 post-treated	7.45 \pm 0.74	8.12 \pm 0.21	4.2 \pm 1.22	1.8 \pm 0.38

Histopathological changes

The figures (9), (10), (11) and (12) showed the histopathological changes in four groups under microscopic examination




Figure 9
the kidney of normal histoarchitecture induced by STZ after 24-day show normal kidney tissues and showed tubes and glomeruli intact. 400 X H & E.




Figure 10
diabetic kidney(G2) of rats the control group, where congestion and severe atrophy of the blood vessels with the destruction of the vascular wall 400 X H & E.




Figure 11

kidney rats in G3 by local EEP that pre-injected with STZ diabetes which stimulates the presence congestion in the shows the breadth light of pipes blood vessels with small areas of red blood cells in the interstitial leak, and also there is a partial improvement in




Figure 12

kidney of rats in G4 by local EEP that induced by STZ show congestion and vascular moderate and of mild to moderate capsules Bowman well-preserved, and shown the structure close to the tissue of normal kidney. 400 XH & E.

histoarchitecture kidney Bowman capsule is retained. 400X H & E.

Microscopic examination showed in the G1 no changes in the microscopic structure of the kidney where it was normal kidneys in the control group. The G2 (diabetic) showed microscopic examination of kidney damage in several inflation and the deterioration of the epithelium of the renal tubules, there vascular congestion, and severe atrophy, with the destruction of the vascular wall, these findings are consistent with^(38,43,44). The G3 pre-treated, which is caused by STZ in rats, where

the microscopic examination, the presence of mild to moderate congestion in the blood vessels, with small areas of red blood cells in the interstitial leak. Bowman capsule is retained. The effect of local EEP, where the microscopic examination showed reduced pathological changes were partially successful in terms of reduction of kidney damage consistent with these results^(38,45). The G4 of rats post-treated, were showed a significant improvement in pathological changes. Microscopic examination showed, the presence of vascular congestion mild, light

and shows the breadth of the pipe and maintain good Bowman capsules, and showed the structure close to normal kidney tissue, this is consistent with^(42,46). In this study, propolis wizard to diabetic nephropathy rats reversed the effective protection and improvement of histopathological changes.

CONCLUSION

Iraqi propolis cannot reduce the genotoxicity induced STZ but has immunological properties through its effect on reducing the concentration of 3-NT and NO[•]. It also has anti-oxidant properties through its effect as anti-hyperglycemic reagent.

REFERENCES

1. Luxton, R. (2010)"Clinical biochemistry" 2nd ed. Vinod Vasishtha for Viva books private limited: Raj press. India: 115,116,218.
2. Ohashi, N.; Urushihara, M.; Satou, R.; and Kobori, H. Glomerular angiotensinogen is induced in mesangial cells in diabetic rats via reactive oxygen species--ERK/JNK pathways: *Hypertens Res.*;33: 1174-1181, 2010.
3. Alter, ML.; Ott, IM.; von Websky, K.; Tsuprykov, O.; Sharkovska, Y.; Relle; Raila; Henze; Klein and Hocher. DPP-4 inhibition on top of angiotensin receptor blockade offers a new therapeutic approach for diabetic nephropathy: *Kidney Blood Press Res.*; 36:119-130, 2012.
4. Alsaman AR; Almashhedy LA; Hadwan MH. Zinc supplementation attenuates lipid peroxidation and increases antiperoxidant activity in seminal plasma of Iraqi asthenospermic men. *Life Sci J*,10:989–997, 2013.
5. Denicola, A.; Souza, JM.; Radi, R. Diffusion of peroxynitrite across erythrocyte membranes. *Proc. Natl. Acad. Sci. U S A*;95: 35: 66–71, 1998.
6. Burney, S.; Caulfield, JL.; Niles, JC; Wishnok, JS. and Tannenbaum SR. The chemistry of DNA damage from nitric oxide and peroxynitrite. *Mutate Res*; 424:37-49, 1999.
7. Hadwan MH; Almashhedy LA; Alsaman AS. Study of the effects of oral zinc supplementation on peroxynitrite levels, arginase activity and NO synthase activity in seminal plasma of Iraqi asthenospermic patients. *Reproductive Biology and Endocrinology*; 12:1, 2014.
8. Jabir FA; Jalil M; Mizil O. A new rout for the evaluation of oxidative stress in diabetic patients by using the activity of some enzymatic antioxidants. *Jornal of Kerbala University*; 5 (2) 23-37, .2007.
9. Mabley, J. and Soriano, FG. Role of nitrosative stress and poly-(ADP-ribose) polymerase activation in diabetic vascular dysfunction. *Curr Vasc. Pharmacol*; 3:247–252, 2005.
10. Pacher, P. and Szabo, C. Role of peroxynitrite in the pathogenesis of cardiovascular complications of diabetes. *Curr Opin Pharmacol*;6:136-141, 2006.
11. Pacher, P.; Obrosova, IG.; Mabley, JG. and Szabo, C. Role of nitrosative stress and peroxynitrite in the pathogenesis of diabetic complications. Emerging new therapeutically strategies. *Curr Med Chem*; 12:267-275, . 2005.
12. Jabir FA; Jalil M; Mizil O. How To Understand The Relationship Between Oxidative Stress, Non Enzymatic Antioxidant And Trace Elements In Diabetic Patients. *Jornal of Kerbala University*; 5 (2): 11-22, 2007.
13. Heilig, C.; Zaloga, C.; Lee, M.; Zhao, X. Riser, B. and Brosius, F. Immunogold localization of high-affinity glucose transporter isoforms in normal rat kidney. *Lab Invest.*; 73:674-84, 1995.
14. Gnudi, L.; Viberti, G.; Raij, L.; Rodriguez, V.; Burt, D. and Cortes, P. GLUT-1overexpression: Link between hemodynamic and metabolic factors in glomerular injury? *Hypertension*; 42:19-24, 2003.
15. Heilig, CW.; Liu, Y.; England, RL; Freytag, SO.; Gilbert, JD. Heilig, KO.; Zhu, M.; Concepcion, LA. and Brosius, FC. D Glucose stimulates mesangial

- cell GLUT1 expression and basal and IGF-I-sensitive glucose uptake in rat mesangial cells: implications for diabetic nephropathy. *Diabetes*; 46:1030-1039, . 1997.
16. Wagner, N.; Morrison, H.; Pagnotta, S.; Michiels, JF.; Schwab, Y.; Tryggvason, K.; Schedl, A. and Wagner, KD. The podocyte protein nephrin is required for cardiac vessel formation. *Human Molecular Genetics*, 20(11): 94,2182, (2011)
 17. Hadwan MH, Almashhedy LA, Alsalman AS. Precise method for the assessment of catalase-like activity in seminal fluids. *Int J Pharm Bio Sci.*, 4(1): 949 – 954, (2013)
 18. Benkovic, V.; Kopjar, N.; Horvat Knezevic, A.; Đikić, D.; Baic, I.; Ramić, S.; Viculin, T.; Knežević, F. and Oršolić, N. Evaluation of radioprotective effects of propolis and quercetin on human white blood cells in vitro. *Biol Pharm Bull.*, 31: 1778-1785, (2008).
 19. Qian, W. L.; Khan, Z.; Watson, D. G. and Fearnley, J. Analysis of sugars in bee pollen and propolis by ligand exchange chromatography in combination with pulsed amperometric detection and mass spectrometry. *Journal of Food Composition and Analysis*, 21(1): 78-83, (2008).
 20. Abo-Salem, O. M.; El-Edel, R. H.; Harisa, G. E. I.; El-Halawany, N. and Ghonaim, M. M. Experimental diabetic nephropathy can be prevented by propolis: effect on metabolic disturbances and renal oxidative parameters. *Pakistan Journal of Pharmaceutical Sciences*. 22(2): 205–210, (2009).
 21. Ito, M. New model of progressive non-insulin-dependent diabetes mellitus in mice induced by Streptozotocin. *Biol Pharmaceut Bull.*, 22: 988-989, (1999).
 22. Ikebukuro, K.; Adachi, Y.; Yamada, Y.; Fujimoto, S.; Seino, Y. and Oyaizu, H. Treatment of Streptozotocin-induced diabetes mellitus by transplantation of islet cells Plus bone Marrow cells via portal vein in rats; *Transplantation*, 73: 512-518, (2002).
 23. Abeer, A. Abd El-Baky. Clinicopathological Effect camellia siensis of Extract on Streptozotocin-Induced Diabetes in Rats. *World Journal of Medical Sciences*, 8 (3): 205-211, (2013).
 24. Cheng, D.; Liang, B. and Li, Y. Antihyperglycemic effect of Ginkgo biloba extract in streptozotocin-induced diabetes in rats. *Bio. M Res. Intern. Journal.*, 1: 1-7, (2013).
 25. Krell, R. Value-added products from bee keeping. *FAO Agricultural Services Bulletin.*, 124: 421, (1996).
 26. Sambrook, J. and Russell, D.W. "Molecular cloning: laboratory manual" 3rd ed. Cold spring Harbor Laboratory Press, *Cold spring Harbor*, N.Y. (2001) 65-72..
 27. Orita, M.; Iwahana, H.; Kanazawa, H.; Hayashi, K. and Sekiya, T. Detection of 200 polymorphisms of human DNA by gel electrophoresis as single-strand conformation 201 polymorphisms. *Proc. Natl. Acad. Sci. USA*, 86: 2766-2770, (1989).
 28. Hortelano, S.; Dewer, B.; Genaro, A.N.; Diaz-Guerra, M.J. and Bosca, L. Nitric oxide is released resulting in regenerating liver after partial hepatectomy. *Hepatology.*, 21: 776-786, (1995).
 29. Fossati, P.; Prencipe, L. and Berti, G. Determination of Uric acid. *Cliniche.*, 1(26): 227-31, (1980).
 30. Jaffe, M. Creatinine Jaffe endpoint methods with deproteinization. *Z. Physiol. Chem.*, 10: 391, (1986).
 31. Login, G. R. and Dvorak, A. M. Methods of microwave fixation for microscopy. A review of research and clinical applications., 25: 1970–1992, (1994).
 32. Gleiberman, A. S.; Fedtsova, N. G. and Rosenfeld, M. G. Tissue interactions in the induction of anterior pituitary: role of the ventral diencephalon, mesenchyme and notochord. *Dev. Biol.*, 213: 340–353, (1999).
 33. Wood, A. E. and Eillis, R. C. "Laboratory histopathology. A complete references". 1st ed. Churchill Livingstone. Edinburgh. (1994) ;10-21.
 34. Tjalve, H. Streptozotocin: distribution, metabolism and mechanisms of action. *Uppsala Med. Sci. Suppl Journal.*, 39: 145-157, (1983).

35. Alejandro, D.; Bolza'n, M. and Bianchi, S. Genotoxicity of streptozotocin. *Elsevier Science B.V. Journal.*, 512: 121-134, (2002).
36. Valadares, B. L.; Graf, U. and Spano, M. A. Inhibitory effect of water extract of propolis on doxorubicin- induced somatic mutation and recombination in *Drosophila melanogaster*. *Food Chem. Toxicol.*, 46:1103-1110, (2008).
37. Nawal, K.J. Effect of Local Extraction Ethanollic Propolis on Some Oxidative Stress Indicators in Rats with Induced Diabetic Nephropathy . (2013) M.Sc. thesis., Department of Chemistry/ College of Education/ University of Al-Qadisiya.
38. El-Sayed, S. M.; Abo- Salman, O. M.; Aly, H. A. and Mansour, A. M. Potential antidiabetic and hypolipidemic effects of propolis extract in streptozotocin-induced diabetic rats. *Pakistan Pharm Sci. Journal.*, 22(2): 74-168, (2009).
39. Antonio, C.; Lisa, Q. and Barbara C. Role of Hyperglycemia in Nitrotyrosine Postprandial Generation: *Diabetic Care*, 25: 8, (2002).
40. Yu-Jung, C. and John Q. Fenofibrate Treatment of Diabetic Rats Reduces Nitrosative Stress, Renal Cyclooxygenase-2 Expression and Enhanced Renal prostaglandin Release: *Valhalla*, (2007).
41. Majida, A.J. and Sawsan, M. Renal effects of propolis and malic acid in Aluminium Exposed Male Rats. *Applied Science Reports Journal.*, 1 (1): 26-30, (2014).
42. Rexlin, S. M., and Rajadurai, M. Effect of Glymin, a Polyherbal Formulation on Lipid Profile and Histopathological Examination in Streptozotocin-Induced Diabetic Rats. *IJPSSDR.*, 1(4): 49-55, (2012).
43. Muhammad, Z.; Syed, N. N.; Masood, A. and Zahid, A. K. Altered Kidney Morphology and Enzymes in Streptozotocin Induced Diabetic Rats. *Int. Morphol Journal.*, 27(3): 783-790, (2009).
44. Mesbah, L.; Kheira, B.; Wided, K. and Mohamed, A. Polyphenolic fractions of Algerian propolis reverses doxorubicin induced acute renal oxidative stress. *Afr. Pharm. Pharmacol Journal*; 4(10): 712-720, (2010).
45. El Agawany, A.; Eiman, M.; Hoda, K. and El Harri, M. Propolis Effect on Rodent Models of Streptozotocin-Induced Diabetic Nephropathy. *Journal of American Science*; 8(12), (2012).