

See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/307537055

Analysis of Volatile Metabolites Released by Staphylococcus Aureus using Gas Chromatography-Mass...

Article in Oriental Journal of Chemistry · August 2016

DOI: 10.13005/ojc/320439

CITATIONS 2		reads 76	reads 76		
3 autho	rs, including:				
	Imad Hadi Hameed University of Babylon 66 PUBLICATIONS 423 CITATIONS SEE PROFILE		Ghaidaa Mohammad University of Al-Qadisiyah/Science college 20 PUBLICATIONS 63 CITATIONS SEE PROFILE		

Some of the authors of this publication are also working on these related projects:



biochemical analysis View project



phytochemical analysis, antibacterial, antifungal View project

All content following this page was uploaded by Imad Hadi Hameed on 05 September 2016.



ORIENTAL JOURNAL OF CHEMISTRY

An International Open Free Access, Peer Reviewed Research Journal

ISSN: 0970-020 X CODEN: OJCHEG 2016, Vol. 32, No. (4): Pg.

www.orientjchem.org

Analysis of Volatile Metabolites Released by *Staphylococcus aureus* using Gas Chromatography-mass Spectrometry and Determination of its Antifungal Activity

HAYFAA HUSSEIN JADDOA¹, IMAD HADI HAMEED*² and GHAIDAA JIHADI MOHAMMED³

¹Department of Biology, Babylon University, Iraq. ²College of Nursing, Babylon University, Iraq. ³College of Science, Al-Qadisiya University, Iraq. *Corresponding author E-mail: imad_dna@yahoo.com

ABSTRACT

Bacterial volatile organic compounds (VOCs) have been considered as sensitive and specific biomarkers for bacterial detection in human specimens and culture media. The possibility of using VOCs markers as one of the largest groups of bacterial metabolites would open a new frontier for developing more efficient techniques in the diagnosis of bacterial infections. The aims of this research were analysis of the bioactive chemical products and evaluation of antibacterial and antifungal activity. Bioactives (chemical compounds often referred to as secondary metabolites) were analyzed using gas chromatography-mass spectroscopy (GC-MS) techniques, then the in vitro antibacterial and antifungal activity of the methanolic extract was evaluated. Thirty five bioactive compounds were identified in the methanolic extract of Staphylococcus aureus. GC-MS analysis of Staphylococcus aureus revealed the existence of the: Hexanoic acid, 2-methyl, 12,15-Octadecadiynoic acid, methyl ester, 1-Aminononadecane ,N-trifluoroacetyl-, N-[3-[N-Aziridyl]propylidene]hexylamine, N-(2,5-Dicyano-3,4-dihydro-2H-pyrrol-2-yl)-acetamide, 3-Cyclohex-3-enyl-propionic acid, 1-Methyl-4-[nitromethyl]-4-piperidinol, 3-Azonia-5-hexyn-1-ol, N,N-dimethyl-O-acetyl-,bromide, 1-Hexadecanol -2-methyl-, 1-Propyl-3,6-diazahomoadamantan, 9-Borabicyclo[3.3.1]nonane, 9-mercapto-, Benzyl methyl ketone, L-Aspartic acid ,N-glycyl-, Aminoacetamide , N-methyl-N-[4-(1-pyrrolidinyl)-2butynyl]-, Tertbutyloxyformamide, N-methyl-N-[4-(1-pyrrolidinyl)-2-butyn, 5,7-Dodecadiyn-1,12diol, Deoxyspergualin, D-Streptamine, O-6-deoxy-á-D-glucopyranosyl-(1-4), dl-Citrulline, N-[3-Diethylaminopropyl]-4-oxo-1,2,3,4,5,6,7,8-octahydroqui, N-Propionyl-D-glucoseamine, Cvstine, 3,4-Dihydrocoumarin ,6-fluoro-4,4-dimethyl-, 4-(2,5-Dihydro-3-methoxyphenyl)butylamine, 3-methoxy-2-(1-methylethyl)-5-(2-methylpropyl)pyrazine, Uric acid, Thiocyanic acid 4-methoxy-2,6-dimethyl-3-pyridyl ester, 12-Dimethylamino-10-oxododecanoic acid, Glycyl-D-asparagine, Actinomycin C2, 12-Octadecenoic acid, methyl ester, 6-Octadecenoic acid, methyl ester, (z)-, 2,5-Piperazinedione , 3,6-bis(2-methylpropyl)-, I-Leucyl-d-leucine and Methyl 12-hydroxy-9-octadecenoate. Gramineae poaceae was very highly active (6.71±0.13) mm. The results of anti-fungal activity produced by Staphylococcus aureus showed that the volatile compounds were highly effective to suppress the growth of Aspergillus terreus. Staphylococcus aureus produce many important secondary metabolites with high biological activities. Based on the significance of employing bioactive compounds in pharmacy to produce drugs for the treatment of many diseases, the purification of compounds produced by Staphylococcus aureus can be useful.

Keywords: Antifungal activity, Staphylococcus aureus, GC-MS, Secondary metabolites.

INTRODUCTION

Staphylococcus aureus is a Grampositive spherical bacterium approximately 1 im in diameter. On a rich medium, S. aureus forms medium size "golden" colonies. One of the important characteristics of S. aureus is its capability to secrete toxins that disrupt membranes of host cells. Cytolytic toxins form β -barrel pores in the cytoplasmic membranes and cause leakage of the cell's content and lysis¹⁻³. Its cells form grape-like clusters, since cell division takes place in more than one plane. It is often found as a commensal associated with skin, skin glands, and mucous membranes, particularly in the nose of healthy individuals⁴. S. aureus secrets several cytolytic toxins, among them alphahemolysin, beta-hemolysin, gamma-hemolysin, leukocidin, and Panton-Valentine leukocidin (PVL)5. It has been estimated that approx. 20-30% of the general population are S. aureus carriers⁶. Staphylococcus aureus can be found in the anterior nares of a great proportion of a healthy human population. S. aureus is one of the main causes of hospital- and community-acquired infections which can result in serious consequences^{7,8}. S. aureus is also able to cause a variety of skin and soft tissue infections and debilitating or even fatal diseases, such as pneumonia, necrotising fasciitis and septicaemia. Nosocomial S. aureus infections affect the bloodstream, skin, soft tissues and lower respiratory tracts. It can also produce toxins which cause toxin-mediated conditions such as toxic shock syndrome or food intoxications. It is a challenge to define which genetic factors determine whether an encounter between a human and S. aureus results in asymptomatic carriage or in clinical disease⁹⁻¹¹. Many bacterial pathogens could lead to life-threatening infections. Accurate and rapid diagnosis is essential for the successful management of these infectious diseases. Traditional bacterial identification methods are time-consuming, require specific techniques and expertise. Other limitations of these techniques such as unaffordability and unavailability of expensive microbiological equipment and delay in the transport of human specimens such as fecal samples in diarrhea patients to the appropriate laboratories remain as main causes of delay in proceeding suitable curative actions in some countries. Therefore, all mentioned reasons have led to unavoidable delay in diagnosis and even death of infected patients¹².

Chemical analysis of bacterial culture includes analysis of bacterial metabolites, bacterial cell wall compositions and fatty acids profiling, have been introduced as bacterial differentiation and detection methods¹³⁻¹⁵. Metabolomics is a fast developing 'omics' that analyzes final metabolites of the cells by means of high throughput analytical technologies such as gas chromatography-mass spectrometry and high performance liquid chromatography-mass spectrometry^{16,17}. Recent advances in ionization technologies allow researchers to perform sensitive qualitative and quantitative analysis of high molecular weight compounds along with the conventional ability of low molecular weight compound analysis in biological experiments¹⁸⁻²¹. The objectives of this research were analysis of the bioactive chemical products and evaluation of antibacterial and antifungal activity.

MATERIALS AND METHODS

Growth conditions and determination of metabolites

S. aureus strain was isolated from bronchitis patients and obtained from Maternity and children hospital. Subcultures were obtained on the Nutrient Agar for 48 hrs. at 22°C. The mixture was incubated at 4°C for 10 min and then shook for 10 min at 130 rpm. Metabolites was separated from the liquid culture and evaporated to dryness with a rotary evaporator at 45°C. The residue was dissolved in 1 ml methanol, filtered through a 0.2 im syringe filter, and stored at 4°C for 24 h before being used for GC-MS. The identification of the components was based on comparison of their mass spectra with those of NIST mass spectral library as well as on comparison of their retention indices either with those of authentic compounds or with literature values²²⁻²⁶.

The studied fungi, Aspergillus niger, Aspergillus terreus, Aspergillus flavus, Aspergillus fumigatus, Candida albicans, Saccharomyces cerevisiae, Penicillium expansum, and Trichoderma viride were isolated and maintained in potato dextrose agar slants. Spores were grown in a liquid culture of potato dextrose broth (PDB) and incubated at 25°C in a shaker for 16 days at 130 rpm. The extraction was performed by adding 25 ml methanol to 100 ml liquid culture in an Erlenmeyer flask after the infiltration of the culture.

Collection and Preparation of Plant Material

In this research, the leaves were dried at room temperature for ten days and when properly dried the leaves were powdered using clean pestle and mortar, and the powdered plant was size reduced with a sieve. The fine powder was then packed in airtight container to avoid the effect of humidity and then stored at room temperature²⁷⁻³¹.

Extraction and Identification of Alkaloids

The powdered leaves (2 g) were boiled in a water bath with 20 ml of 5% sulphuric acid in 50% ethanol. The mixture was cooled and filtered. A portion was reserved. Another portion of the filtrate was put in 100 ml of separating funnel and the solution was made alkaline by adding two drops of concentrated ammonia solution. Equal volume of chloroform was added and shaken gently to allow the layer to separate. The lower chloroform layer was run off into a second separating funnel³²⁻³³. The ammoniacal layer was reserved. The chloroform layer was extracted with two quantities each of 5 ml of dilute sulphuric acid. The various extracts were then used for the following test:

Wagner's test: To the filtrate in tube III, 1 ml of wagner's reagent was added drop by drop. Formation of a reddish-brown precipitate indicates the presence of alkaloids.

Dragendoff's test: To the filtrate in test tube II, 1 ml of dragendoff's reagent was added drop by drop. Formation of a reddish-brown precipitate indicates the presence of alkaloids.

Mayer's test: To the filtrate in test tube I, 1 ml of mayer's reagent was added drop by drop. Formation of a greenish coloured or cream precipitate indicates the presence of alkaloids.

Spectral analysis of bioactive chemical compounds using gas chromatography-mass spectrometry (GC/MS)

Analysis was conducted using GC-MS (Agilent 789 A) equipped with a DB-5MS column ($30 \text{ m} \times 0.25 \text{ mm i.d.}$, 0.25 um film thickness, J&W Scientific, Folsom, CA). The oven temperature was programmed as for the previous analysis. Helium was used as the carrier gas at the rate of 1.0 mL/

min. Effluent of the GC column was introduced directly into the source of the MS via a transfer line (250°C). Ionization voltage was 70 eV and ion source temperature was 230°C. Scan range was 41- 450 amu. The components were identified by comparing their retention times to those of authentic samples of Wiley Mass Spectral data base Library³⁴⁻³⁸.

Determination of antibacterial activity

Five-millimeter diameter wells were cut from the agar using a sterile cork-borer, and 25 il of the samples solutions (Piper nigrum, Zingiber officinale, Gramineae poaceae, Nerium olender, Ricinus communis, Datura stramonium, Linum usitatissimum, Anastatica hierochuntica, Cassia angustifolia, Euphorbia lathyrus, Rosmarinus oficinalis, Mentha viridis, Artemisia annua, Quercus infectoria, Citrullus colocynthis, Althaea rosea, Coriandrum sativum, Origanum vulgare, Urtica dioica, Equisetum arvense, Foeniculum vulgare, Nigella sativa, Ocimum basilicum, Punica granatum, Punica granatum and Cinnamomum zeylanicum) were delivered into the wells. The plates were incubated for 48 h at room temperature. Antimicrobial activity was evaluated by measuring the zone of inhibition against the test microorganisms. Methanol was used as solvent control. The tests were carried out in triplicate. The antifungal activity was evaluated by measuring the inhibition-zone diameter observed after 48 h of incubation³⁹⁻⁴⁹.

Data analysis

All the measurements were replicated three times for each assay and the results are presented as mean \pm SD and mean \pm SE. IBM SPSS 20 version statistical software package was used for statistical analysis of percentage inhibition and disease incidence and disease severity in each case.

RESULTS AND DISCUSSION

Gas chromatography and mass spectroscopy analysis of compounds was carried out in methanolic extract of *S. aureus*, shown in Table 1. The GC-MS chromatogram of the thirty one peaks of the compounds detected was shown in Figure 1. Peaks were determined to be Hexanoic acid, 2-methyl, 12,15-Octadecadiynoic acid, methyl ester, 1-Aminononadecane ,N-trifluoroacetyl-, N-[3-[N-Aziridyl]propylidene]hexylamine, N-(2,5-

methanolic extract of Staphylococcus aureus OH 12,15-Octadecadiynoic acid , methyl 1-Aminononadecane ,N-trifluoroacetyl-Hexanoic acid, 2-methylester RT=3.453, Mw=290.22458 RT=3.149, Mw=130.09938 RT=3.619, Mw=379.3062, Pharmacological activity: Pharmacological activity: Pharmacological activity: anti-microbial activity anti-inflammatory compound anti-malarial and anti-convulsant N-[3-[N-Aziridyl]propylidene] N-(2,5-Dicyano-3,4-dihydro-2H-3-Cyclohex-3-enyl-propionic acid hexylamine, RT=3.745, Mw=182.178299 pyrrol-2-yl)-acetamide, RT=3.922, RT=4.265, Mw=154.09938, Mw=176.069811, Pharmacological Pharmacological activity: anti-viral Pharmacological activity: activity: Antibacterial and antifungal anti-thrombotic activity activities OH 0/ 1-Methyl-4-[nitromethyl]-4-piperidinol 3-Azonia-5-hexyn-1-ol, N,N-dimethyl-1-Hexadecanol -2-methyl-RT=4.398, Mw=174.100442 O-acetyl-, bromide, RT=4.448 RT=4.569, Mw=256.276615, Pharmacological activity: Mw=170.118104, Pharmacological Pharmacological activity: activity: antifungal, molluscicides and antimicrobial, antioxidant Anti-inflammatory activity antileishmanial 1-Propyl-3,6-diazahomoadamantan 9-Borabicyclo[3.3.1]nonane, Benzyl methyl ketone RT=4.734, Mw=210.173213 9-mercapto- RT=5.158 RT=5.307, Mw=134.073165 Mw=154.098752, Pharmacological Pharmacological activity: Pharmacological activity: antibacterial and antifungal, antiviral, anti-inflammatory activities activity: Unknown

antiparasitic, anti-tubercular

Table 1: Bioactive chemical compounds identified in



L-Aspartic acid ,N-glycyl-RT=5.530, Mw=190.058971 Pharmacological activity: anti-inflammatory



Aminoacetamide , N-methyl-N-[4-(1pyrrolidinyl)-2-butynyl]-RT=7.166 Mw=209.152812, Pharmacological activity: Anti-tumor



Tertbutyloxyformamide, N-methyl-N-[4-(1-pyrrolidinyl)-2-butyn RT=7.664, Mw=252.183778 Pharmacological activity: anti-thrombotic activities

HO OH

5,7-Dodecadiyn-1,12-diol RT=8.025, Mw=194.13068 Pharmacological activity: antitumor and *anti*-obesity activities



Deoxyspergualin RT=8.168, Mw=387.295788 Pharmacological activity: *anti*- tumor *activity*



D-Streptamine , O-6-deoxy-á-Dglucopyranosyl-(1-4) RT=8.093 Mw=482.258793, Pharmacological activity: *Anti*-Bacteria



dl-Citrulline RT=8.374, Mw=175.095692 Pharmacological activity: *anti*-asthenic *effect*



N-[3-Diethylaminopropyl]-4-oxo-1,2,3,4,5,6,7,8-octahydroqui RT=10.096, Mw=264.220163 Pharmacological activity: Unknown



N-Propionyl-D-glucoseamine RT=10.399, Mw=235.105587 Pharmacological activity: Anti-tumor properties

NH₂ NH₂

Cystine, RT=10.582 Mw=240.023849 Pharmacological activity: antithrombotic agents or tumour growth inhibitors



3,4-Dihydrocoumarin ,6-fluoro-4,4dimethyl-RT=11.069 Mw=194.074307, Pharmacological activity: *anti*-ulcer activity



4-(2,5-Dihydro-3-methoxyphenyl) butylamine, RT=11.395 Mw=181.146665, Pharmacological activity: *anti*-inflammatory activities



Thiocyanic acid 4-methoxy-2,6dimethyl-3-pyridyl ester, RT=14.033 Mw=194.051384, Pharmacological activity: *anti*-inflammatory activity



Uric acid, RT=12.659 Mw=168.02834



3-methoxy-2-(1-methylethyl)-5-(2methylpropyl)pyrazine, RT=12.196 Mw=208.157563, Pharmacological activity: *anti*-tumor *activity*

OH NH₂ HN NH₂

Glycyl-D-asparagine, RT=14.691 Mw=189.074956, Pharmacological activity: analgesic activity

0

12-Octadecenoic acid, methyl ester RT=16.224, Mw=296.27153, Pharmacological activity: antimicrobial







I-LeucyI-d-leucine, RT=17.678 Mw=244.178693, Pharmacological activity: antibacterial *activity*



6-Octadecenoic acid , methyl ester ,(z)-RT=16.636, 296.27153, Pharmacological activity: have strong analgesic, *anti*inflammatory and antipyretic

2,5-Piperazinedione , 3,6-bis(2-ethylpropyl)-, RT=17.243 Mw=226.168128, Pharmacological activity: Unknown

0H Nethyl 12-hydroxy-9-octadecenoate

Methyl 12-hydroxy-9-octadecenoate RT=17.924, Mw=312.266445 Pharmacological activity: biological activities including antimicrobial and anticancer

6

Dicyano-3,4-dihydro-2H-pyrrol-2-yl)-acetamide, 3-Cyclohex-3-enyl-propionic acid, 1-Methyl-4-[nitromethyl]-4-piperidinol, 3-Azonia-5-hexyn-1-ol, N,N-dimethyl-O-acetyl-,bromide, 1-Hexadecanol -2-methyl-, 1-Propyl-3,6-diazahomoadamantan, 9-Borabicyclo[3.3.1]nonane, 9-mercapto-, Benzyl methyl ketone, L-Aspartic acid ,N-glycyl-, Aminoacetamide , N-methyl-N-[4-(1-pyrrolidinyl)-2-butynyl]-, Tertbutyloxyformamide , N-methyl-N-[4-(1-pyrrolidinyl)-2-butyn, 5,7-Dodecadiyn-1,12-diol, Deoxyspergualin, D-Streptamine , O-6-deoxy-α-D-glucopyranosyl-(1-4), dl-Citrulline,

Fungi	Antibiotics / Staphylococcus aureus metabolite products				
	Metabolite products of <i>Staphylococcus</i> <i>aureus</i>	Amphotericin B	Fluconazol	Miconazole nitrate	
Aspergillus niger	5.06±0.21 ª	1.02±0.11	1.94±0.09	1.07±0.10	
Aspergillus terreus	6.08±0.29	2.06±0.16	1.02±0.16	2.01±0.12	
Aspergillus flavus	5.06±0.16	1.04±0.09	2.69±0.24	1.87±0.14	
Aspergillus fumigatus	5.67±0.20	0.97±0.10	3.01±0.14	1.01±0.13	
Candida albicans	5.01±0.11	1.48±0.12	1.79.±0.13	1.05±0.17	
Saccharomyces cerevisiae	3.03±0.17	1.02±0.12	1.03±0.10	2.13±0.12	
Penicillium expansum	2.68±0.14	2.03±0.20	3.90±0.15	0.86±0.11	
Trichoderma viride	3.35±0.20	2.06±0.11	1.37±0.12	1.32±0.12	
Trichoderma horzianum	3.02±0.19	0.39±0.01	2.05±0.13	1.97±0.16	

Table 2: Antifungal activity of Staphylococcus aureus metabolite products

^a The values (average of triplicate) are diameter of zone of inhibition at 100 mg/mL methanolic extract and 30 jg/mL of (Amphotericin B; Fluconazol and Miconazole nitrate)



Fig. 1: GC-MS chromatogram of methanolic extract of *Staphylococcus aureus* chromatogram of methanolic extract of *Staphylococcus aureus*.

N-[3-Diethylaminopropyl]-4-oxo-1,2,3,4,5,6,7,8octahydroqui, N-Propionyl-D-glucoseamine, Cystine, 3,4-Dihydrocoumarin ,6-fluoro-4,4-dimethyl-, 4-(2,5-Dihydro-3-methoxyphenyl)butylamine, 3-methoxy-2-(1-methylethyl)-5-(2-methylpropyl)pyrazine, Uric acid, Thiocyanic acid 4-methoxy-2,6-dimethyl-3pyridyl ester, 12-Dimethylamino-10-oxododecanoic acid, Glycyl-D-asparagine, Actinomycin C2, 12-Octadecenoic acid , methyl ester, 6-Octadecenoic acid , methyl ester ,(z)-, 2,5-Piperazinedione , 3,6bis(2-methylpropyl)-, I-Leucyl-d-leucine and Methyl 12-hydroxy-9-octadecenoate.

Antibacterial and antifungal activity

The results of anti-fungal activity produced by *Staphylococcus aureus* showed that the volatile compounds were highly effective to suppress the growth of *Aspergillus terreus*. *Staphylococcus aureus* produce many important secondary metabolites with high biological activities. Based on the significance of employing bioactive compounds in pharmacy to produce drugs for the treatment of many diseases, the purification of compounds produced by *Staphylococcus aureus* can be useful. Maximum zone formation against *Aspergillus terreus* (6.08±0.29) mm, Table 2. In agar well diffusion

Table 3: Zone of inhibition (mm) of test different bioactive
compounds and standard antibiotics of medicinal plants to
Staphylococcus aureus

S. No.	Plant	Zone of inhibition (mm)
1.	Piper nigrum (Crude)	5.35±0.19
2.	Zingiber officinale (Crude)	4.22±0.20
3.	Gramineae poaceae (Crude)	6.71±0.13
4.	Nerium olender (Alkaloids)	3.36±0.20
5.	Ricinus communis (Alkaloids)	1.86±0.19
6.	Datura stramonium(Alkaloids)	2.52±0.20
7.	Linum usitatissimum (Crude)	3.88±0.14
8.	Anastatica hierochuntica (Crude)	4.17±0.18
9.	Linum usitatissimum (Crude)	3.43±0.13
10.	Cassia angustifolia (Crude)	5.51±0.25
11.	Euphorbia lathyrus (Crude)	4.69±0.27
12.	Rosmarinus oficinalis (Crude)	4.35±0.15
13.	Mentha viridis (Crude)	3.29±0.14
14.	Artemisia annua (Crude)	2.46±0.28
15.	Quercus infectoria (Crude)	5.19±0.10
16.	Citrullus colocynthis (Crude)	3.00±0.20
17.	Althaea rosea (Crude)	4.70±0.15
18.	Coriandrum sativum (Crude)	4.09±0.20
19.	Melia azedarach (Crude)	2.76±0.24
20.	Origanum vulgare (Crude)	5.15±0.26
21.	Urtica dioica (Crude)	2.11±0.13
22.	Equisetum arvense (Crude)	4.00±0.12
23.	Foeniculum vulgare (Crude)	2.25±0.15
24.	Nigella sativa (Crude)	3.27±0.24
25.	Ocimum basilicum (Crude)	3.18±0.15
26.	Punica granatum (Crude)	5.36±0.20
27.	Cinnamomum zeylanicum (Crude)	3.01±0.18
28.	Streptomycin	2.17±0.10
29.	Rifambin	1.04±0.11
30.	Control	00.00

method the selected medicinal plants (Piper nigrum, Zingiber officinale, Gramineae poaceae, Nerium olender, Ricinus communis, Datura stramonium, Linum usitatissimum, Anastatica hierochuntica, Cassia angustifolia, Euphorbia lathyrus, Rosmarinus oficinalis, Mentha viridis, Artemisia annua, Quercus infectoria, Citrullus colocynthis, Althaea rosea, Coriandrum sativum, Origanum vulgare, Urtica dioica, Equisetum arvense, Foeniculum vulgare, Nigella sativa, Ocimum basilicum, Punica granatum, Punica granatum and Cinnamomum zeylanicum) were effective against Staphylococcus aureus, Table 3. Gramineae poaceae was very highly active (6.71±0.13) mm against Staphylococcus aureus. Staphylococcus aureus was found to be sensitive to all test medicinal plants and mostly comparable to the standard reference antifungal drug Amphotericin B and fluconazole to some extent. Recently, it was demonstrated that volatile organic compounds (VOCs) of bacteria such as terpenoids, phenylpropanoids and fatty acid derivatives can influence the growth of some fungi and, in general, the inter- and intra-organismic communication signals.

CONCLUSION

Thirty five bioactive chemical constituents have been identified from methanolic extract of the *Staphylococcus aureus* by gas chromatogram mass spectrometry (GC-MS). In vitro antifungal and antibacterial evaluation of secondary metabolite products of *Staphylococcus aureus* forms a primary platform for further phytochemical and pharmacological investigation for the development of new potential antimicrobial compounds.

ACKNOWLGEMENTS

I would like to express my deep appreciation to head of department Prof. Dr. Ali Al-Marzooqi and Dean of the Science College Prof. Dr. Inas Al-Rubae for their support and encouragement.

REFERENCES

- 1. Foster, T.J. *Nat Rev Microbiol.* **2005**, *3*, 948–958.
- Mohammed, G.J.; Kadhim, M.J.; Hussein, H.M. International Journal of Pharmacognosy and Phytochemical Research. 2016, 8(6), 889-905.
- Filipiak, W.; Sponring, A.; Baur, M.M.; Filipiak, A.; Ager, C. *BMC Microbiol.* 2012, *12*, 113.
- Buszewski, B.; Ulanowska, A.; Ligor, T.; Jackowski, M.; Klodzinska, E. Journal of Chromatography B: Biomedical Sciences and Applications. 2008, 868, 88-94.
- 5. Kaneko, J.; Kamio, Y. *Biosci Biotechnol Biochem.* **2004**, *68*, 981–1003.
- Kunze, N.; Göpel, J.; Kuhns, M.; Jünger, M.; Quintel, M. *Applied microbiology and biotechnology*. 2013, *97*, 3665-3676.
- Schulz, S.; Dickschat, J.S. Nat Prod Rep. 2007, 24, 814-842.
- 8. Zhu, J.; Bean, H.D.; Kuo, Y.M.; Hill, J.E. *J Clin Microbiol.* **2010**, *48*, 4426-4431.
- Becker, K.; Friedrich, A.W.; Lubritz, G.; Weilert, M.; Peters, G.; Von Eiff, C. J *Clin Microbiol.* 2003, *41*, 1434–1439.
- 10. Fueyo, J.M., Mendoza, M.C., Martín, M.C.

Microbes Infect, 2005, 7, 187–194.

- Belkum, A.; Melles, D.C.; Nouwen, J., Leeuwen, W.B.; Wamel, W., Vos, M.C., Wertheim, H.F.; Verbrugh, H.A. Infect Genet Evol, 2009, 9, 32–47.
- 12. Probert, C.; Jones, P.; Ratcliffe, N. *GUT*, **2004**, *53*, 58-61.
- Ehrhardt, C.J.; Chu, V.; Brown, T.; Simmons, T.L.; Swan, B.K. *Applied and Environmental Microbiology*, **2010**, *76*, 1902-1912.
- 14. Li, Y., Wu, S., Wang, L.; Li, Y.; Shi, F. *J Sci Food Agric.* **2010**, *90*, 1380-1383.
- 15. Pagans, E.; Font, X.; Sánchez, A. Journal of hazardous materials. 2006, 131, 179-186.
- Wang, D.; Ding, X.; Rather, P.N. Journal of Bacteriology. 2001, 183, 4210-4216.
- Savelev, S.U.; Perry, J.D.; Bourke, S.J.; Jary, H.; Taylor, R. *Lett Appl Microbiol.* 2011, *52*, 610-613.
- 18. Glish, G.L.; Vachet, R.W. *Nature Reviews Drug Discovery*. **2003**, *2*, 140-150.
- Ulanowska, A.; Kowalkowski, T.; Hrynkiewicz, K.; Jackowski, M.; Buszewski, B. *Biomed Chromatogr*, 2011, *25*, 391-397.
- 20. Garner, C.E.; Smith, S.; Elviss, N.C.;

Humphrey, T.J.; White, P. Biomarkers. **2008**, *13*, 413-421.

- 21. Davies, J.C. Paediatr Respir Rev. 2002, 3, 128-134.
- 22. Liu, G.Y.; Essex, A.; Buchanan, J.T.; Datta, V.; Hoffman, H.M.; Bastian, J.F. *J Exp Med* **2005**, *202*, 209–215.
- Altameme, H.J.; Hameed, I.H.; Abu-Serag, N.A. *Malaysian Applied Biology*, **2015b**, 44(4), 47–58.
- 24. Mohammed, G.J.; Al-Jassani, M.J.; Hameed, I.H. International Journal of Pharmacognosy and Phytochemical Research, **2016**, *8*(3), 480-494.
- Hussein, H.J.; Hadi, M.Y.; Hameed, I.H. Journal of Pharmacognosy and Phytotherapy, 2016, 8(3), 60-89.
- Shareef, H.K.; Muhammed, H.J.; Hussein, H.M.; Hameed, I.H. Oriental Journal of Chemistry, 2016, 32(2), 20-40.
- Hamza, L.F.; Kamal, S.A.; Hameed, I.H. Journal of Pharmacognosy and Phytotherapy, 2015, 7(9), 194-220.
- Al-Jassaci, M.J.; Mohammed, G.J.; Hameed, I.H. International Journal of Pharmaceutical and Clinical Research, 2016, 8(5), 304-315.10.
- Hameed, I.H.; Ibraheam, I.A.; Kadhim, H.J. Journal of Pharmacognosy and Phytotherapy, 2015c, 7(6), 90-106.
- Al-Marzoqi, A.H.; Hadi, M.Y.; Hameed, I.H. Journal of Pharmacognosy and Phytotherapy, 2016, 8(2), 25-48.
- Hadi, M.Y.; Mohammed, G.J.; Hameed, I.H. Journal of Pharmacognosy and Phytotherapy, 2016, 8(2), 8-24.
- Kadhim, M.J.; Mohammed, G.J.; Hameed, I.H. Oriental Journal of Chemistry. 2016; 32(2), 10-30.
- Jasim, H.; Hussein, A.O.; Hameed, I.H.; Kareem, M.A. *Journal of Pharmacognosy* and Phytotherapy, **2015**, 7(4), 57-72.
- Altameme, H.J.; Hameed, I.H.; Idan, S.A.; Hadi, M.Y. Journal of Pharmacognosy and Phytotherapy. 2015c, 7(9), 222-237.
- 35. Hussein, A.O.; Mohammed, G.J.; Hadi, M.Y.;

Hameed, I.H. *Journal of Pharmacognosy and Phytotherapy*, **2016**, *8*(3), 49-59.

- Hameed, I.H.; Altameme, H.J.; Idan, S.A. Research Journal of Pharmaceutical, Biological and Chemical Sciences. 2016, 7(2), 1843-1868.
- Ratjen, F. Current opinion in pulmonary medicine, 2006, 12, 428-432.
- Shestivska, V., Španel, P.; Dryahina, K.; Sovová, K.; Smith, D. Journal of Applied Microbiology. 2012, 113, 701-713.
- Zechman, J.M.; Aldinger, S.; Labows, J.N. Journal of Chromatography B: Biomedical Sciences and Applications, 1986, 377, 49-57.
- Hameed, I.H.; Hussein, H.J.; Kareem, M.A.; Hamad, N.S. *Journal of Pharmacognosy and Phytotherapy*, **2015b**, *7(7)*, 107-125.
- Sosa, A.A.; Bagi, S.H.; Hameed, I.H. International Journal of Pharmacognosy and Phytochemical Research, 2016, 8(5), 109-126 ().
- 42. Hussein, H.M.; Hameed, I.H.; Ibraheem, O.A. International Journal of Pharmacognosy and Phytochemical Research, **2016**, *8(3)*, 369-385.
- Mohammed, G.J.; Omran, A.M.; Hussein, H.M. International Journal of Pharmacognosy and Phytochemical Research. 2016, 8(6), 977-996.
- 44. Kadhim, M.J.; Sosa, A.A.; Hameed, I.H. International Journal of Pharmacognosy and Phytochemical Research, **2016**, *8(6)*, 127-146.
- Altameme, H.J.; Hadi, M.Y.; Hameed, I.H. Journal of Pharmacognosy and Phytotherapy. 2015a, 7(10), 238-252.
- Hameed, I.H.; Hamza, L.F.; Kamal, S.A. Journal of Pharmacognosy and Phytotherapy. 2015a, 7(8), 132-163.
- Kaur, R.; Kaur, H. Oriental Journal of Chemistry. 2015, 31(1), 597-600.
- 48. Chandrashekharaiah, K.S. Oriental Journal of Chemistry. 2013, 29(3), 1061-1070.
- 49. Kadhim, M.J.; Mohammed, G.J.; Hameed, I.H. Oriental Journal of Chemistry. **2016**, *32(3)*.