

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/304945495>

# Antibacterial and phytochemical analysis of piper nigrum using gas chromatography – mass spectrum and...

Article · June 2016

---

CITATIONS

8

READS

30

3 authors, including:



Ghaidaa Mohammad

University of Al-Qadisiyah/Science college

20 PUBLICATIONS 63 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Phytochemical analysis, heavy metals [View project](#)



phytochemical analysis,antibacterial,antifungal [View project](#)

# Antibacterial and Phytochemical Analysis of *Piper nigrum* using Gas Chromatography – Mass Spectrum and Fourier-Transform Infrared Spectroscopy

Ghaidaa Jihadi Mohammed<sup>1</sup>, Aseel Mohammad Omran<sup>2</sup>, Haider Mashkoor Hussein<sup>1</sup>

<sup>1</sup>College of Science, Al-Qadisiya University, Iraq

<sup>2</sup>College of Pharmacy, Babylon University, Iraq

Available Online: 9<sup>th</sup> June, 2016

## ABSTRACT

The aims of the study were to investigate the presence of phytochemical compounds from the fruits of *Piper nigrum*, using methanolic extraction and report the main functional components by using fourier-transform infrared spectroscopy. The phytochemical compound screened by GC-MS method. A total of 55 bioactive phytochemical compounds were identified in the methanolic extract of *P. nigrum*. The identification of phytochemical compounds is based on the peak area, retention time molecular weight, molecular formula, chemical structure, MS Fragment ions and pharmacological actions. GC-MS analysis of *P. nigrum* revealed the existence of the Propanedioic acid , dimethyl ester, Bicyclo[3.1.1]heptane,6,6-dimethyl-2-methylene-, (1S), 3-Carene, Cyclohexene, 1-methyl-5-(1-methylethenyl)-,(R), 1,6-Octadien-3-ol,3,7-dimethyl, 2-Methyl-1-ethylpyrrolidine, 2-Isopropenyl-5-methylhex-4-enal, L- $\alpha$ -Terpineol, (R)-lavandulyl acetate, Pyrrolizin-1,7-dione-6-carboxylic acid , methyl(ester), 7-epi-cis-sesquisabinene hydrate, Phenol, 2-methoxy-4-(1-propenyl)-(Z), Eugenol, Alfa.Copaene, Naphthalene,1,2,3,5,6,7,8,8a-octahydro-1,8a-dimethyl-7-(1-methyl), Epiglobulol, Caryophyllene, 1,4,7-Cycloundecatriene, 1,5,9,9-tetramethyl-,Z,Z,Z,  $\alpha$ -ylangene,  $\beta$ -copaene, Cedran-diol,8S,13, Isocalamendiol, Cinnami acid , 4-hydroxy -3-methoxy-, {5-hydroxy-2-hydroxymethyl, (-)-Spathulenol, 1-Heptatriacotanol, Desacetylanquidine, 5-Isopropyl-2,8-dimethyl-9-oxatricyclo[4.4.0.0.(2,8)]decan-7-one, Estra-1,3,5(10)-trien-17 $\beta$ -ol, Trans-1,2-Diaminocyclohexane-N,N,N',N'-tetraacetic acid, Phytol, Piperidine,1-(1-oxo-3-phenyl-2-propenyl)-, Eicosanoic acid , 2-(acetoxyo-0-1-[(acetoxy)methyl]ethyl ester, 2,5,5,8a-Tetramethyl-6,7,8,8a-tetrahydro-5H-chromen-8-ol, Z-5-methyl-6-heneicosen-11-one, 2H-1,2-Benzoxazine-3-carbonitrile,2- cyclohexyloctahydro-4a,8a-d, Indoxazin-4-one ,4,5,6,7-tetrahydro-3-undecyl, 9,10-Secococholesta-5,7,10(19)-triene-3,24,25-triol,(3 $\beta$ ,5Z,7E), 3-Oxo-10(14)-epoxyguai-11(13)-en-6,12-olide, 7-[2-(Ethoxycarbonyl)-3 $\alpha$ ,5 $\beta$ -dimethoxycyclopentyl -1]-heptanoic acid, 2H-Benzo[f]oxireno[2,3-E]benzofuran-8,(9H)-one,9-[(1,3-benzodio, Nalorphine, 2-Cyclohexen -3-ol-1-one , 2-[1-iminotetradecyl]-, Piperine, Fenretinide, 11-Dehydrocorticosterone, 5H-Cyclopropano[3,4]benz[1,2-e]azulen-5-one,1,1a,1b,4,4a,7a,7b, 17a-Ethyl-3 $\beta$ -methoxy-17a-aza-D-homoandrost-5-ene-17-one, Bufo-20,22-dienolide m,14,15-epoxy-3,11-dihydroxy-, (3 $\beta$ ,5 $\beta$ ,11 $\alpha$ ,15, 9-Desoxo-9-x-acetoxy-3,8,12-tri-O-acetylngol, Retinal ,9-cis-, 6- $\beta$ -Naltrexol, Piperine, Ursodeoxycholic acid, 5 $\alpha$ -Cholan-24-oic acid , 12 $\alpha$ -hydroxy-3,7-dioxo-,methyl ester and Stigmasterol. The FTIR analysis of *P. nigrum* leaves proved the presence of Alkenes, Aliphatic fluoro compounds, Alcohols, Ethers, Carboxlic acids, Esters and Nitro Compounds. Methanolic extract of bioactive compounds of *P. nigrum* was assayed for *in vitro* antibacterial activity against *Escherichia coli*, *Pseudomonas aerogenosa*, *Proteus mirabilis*, *Staphylococcus aureus* and *Klebsiella pneumonia* by using the diffusion method in agar. The zone of inhibition were compared with different standard antibiotics. The diameters of inhibition zones ranged from 5.00 $\pm$ 0.16 to 0.40 $\pm$ 0.12 mm for all treatments.

**Keywords:** FT-IR, GC-MS analysis, *Piper nigrum*.











## INTRODUCTION

*Piper nigrum* L. is a flowering vine in the family of piperaceae, therefore an important medicinal plant is used in traditional medicine in Asia and Pacific islands especially in Indian medicine<sup>1</sup>. Pepper aroma and flavor due to their chemical substances especially the volatile oil. Black pepper oil has medicinal values. It can be used to help in treatment of pain relief, rheumatism, chills, flu, colds exhaustion, muscular aches, physical and emotional coldness, fever as nerve tonic and to increase circulation<sup>2</sup>.











<sup>5</sup>. The main pungent principle in the green berries of pepper *P. nigrum* L. is piperine. Generally the piperine content of black or white peppercorns lies within the range of 3-8 g/100 g, whereas the content of minor alkaloids piperyline and piperettine have been estimated as 0.2-0.3 and 0.2-1.6 g/100g respectively. The bioavailabilityenhancing property of piperine indicates its potential to be used as an adjuvant with therapeutic drugs in chronic ailments, to reduce the effective dose of the drug and, hence,

Table 1: Major phytochemical compounds identified in methanolic extract of *Piper nigrum*.

S. No.	Phytochemical compound	RT (min)	Formula	Mol. Wt.	Exact Mass	Chemical structure	MS Fragment-ions	P'ecological actions
1.	Propanedioic acid , dimethyl ester	3.150	C <sub>5</sub> H <sub>8</sub> O <sub>4</sub>	132	132.042		59,74,101,132	Anti-tumor and antioxidant activity
2.	Bicyclo[3.1.1] heptane,6,6-dimethyl-2-methylene-,(1S)-	3.339	C <sub>10</sub> H <sub>16</sub>	136	136.1252		53,69,79,936	Anti-Helicobacter pylori activity and Anti-Candida activity
3.	3-Carene	3.693	C <sub>10</sub> H <sub>16</sub>	136	136.1252		53,67,79,936	Several biological activities such as anti-inflammatory
4.	Cyclohexene , 1-methyl-5-(1-methylethethyl)-,(R)-	3.997	C <sub>10</sub> H <sub>16</sub>	136	136.1252		53,68,79,936	Anti-microbial agents
5.	1,6-Octadien-3-ol,3,7-dimethyl-	5.067	C <sub>10</sub> H <sub>18</sub> O	154	154.135765		55,71,80,834	Pharmacologic effects such as: anti-inflammation
6.	2-Methyl-1-ethylpyrrolidine	5.696	C <sub>7</sub> H <sub>15</sub> N	113	113.1204495		56,69,84,98113	Anti-tumor activity.
7.	2-Isopropenyl-5-methylhex-4-enal	6.108	C <sub>10</sub> H <sub>16</sub> O	152	152.120115		69,84,95,109,123,137	Antihyperglycemic and anticariogenic
8.	L- $\alpha$ -Terpineol	6.303	C <sub>10</sub> H <sub>18</sub> O	154	154.135765		55,59,67,819,107,121,136	Anti-inflammatory
9.	(R)-lavandulyl acetate	6.629	C <sub>12</sub> H <sub>20</sub> O <sub>2</sub>	196	196.14633		53,69,80,936,154	Anti-inflammatory
10.	Pyrrolizin-1,7-dione-6-carboxylic acid , methyl(ester)	7.562	C <sub>9</sub> H <sub>11</sub> NO <sub>4</sub>	197	197.068808		55,69,84,987	Anti-Viral and Anti-Tumor Activity
11.	7-epi-cis-sesquisabinene hydrate	8.357	C <sub>15</sub> H <sub>26</sub> O	222	222.198365		55,69,82,93105,119,133,147,161,175,189,204,222	Antioxidant effects and anti-mutagenic action

12.	Phenol , 2-methoxy-4-(1-propenyl)-(Z)-	8.494	C <sub>10</sub> H <sub>12</sub> O <sub>2</sub>	164 73	164.083		51,55,65,77 ,91,103,115 ,121,131,14 9,164	Anti-inflammatory effects
13.	Eugenol	8.523	C <sub>10</sub> H <sub>12</sub> O <sub>2</sub>	164 73	164.083		51,55,65,77 ,91,103,115 ,121,131,13 7,149,164	Anti-Inflammatory agents and antioxidants
14.	Alfa.Copaene	8.740	C <sub>15</sub> H <sub>24</sub>	204 8	204.187		55,69,77,91 ,105,119,13 3,147,161,1 75,189,204	Analgesic and anti-inflammatory activities
15.	Naphthalene,1,2,3,5,6,7,8,8a-octahydro-1,8a-dimethyl-7-(1-methyl	8.958	C <sub>15</sub> H <sub>24</sub>	204 8	204.187		55,67,79,10 7,119,133,1 61,175,189, 204	Anti-bacterial
16.	Epiglobulol	9.181	C <sub>15</sub> H <sub>26</sub> O	222 365	222.198		55,69,82,93 ,109,121,14 7,161,204,2 22	Antioxidant activity, antimicrobial, anti-inflammatory
17.	Caryophyllene	9.456	C <sub>15</sub> H <sub>24</sub>	204 8	204.187		55,69,79,93 ,105,120,13 3,147,161,1 75,189,204	Several biological activities are attributed to beta-caryophyllene, such as anti-inflammatory, antibiotic, antioxidant, anticarcinogenic and local anaesthetic
18.	1,4,7-Cycloundecatriene, 1,5,9,9-tetramethyl-,Z,Z,Z-	9.810	C <sub>15</sub> H <sub>24</sub>	204 8	204.187		55,67,80,93 ,107,121,13 6,147,161,1 75,189,204	Anti-aging, anti-hyperlipidemia and antimicrobial activities
19.	$\alpha$ - ylangene	10.00	C <sub>15</sub> H <sub>24</sub>	204 5	204.187		55,67,77,93 ,105,119,13 3,147,161,1 89,204	Anti-inflammatory properties
20.	$\beta$ -copaene	10.08	C <sub>15</sub> H <sub>24</sub>	204 5	204.187		55,79,105,1 61,204	Anti-neuroinflammatory and neuroprotective
21.	Cedran-diol,8S,13-	10.46	C <sub>15</sub> H <sub>26</sub> O <sub>2</sub>	238 28	238.193		55,74,93,10 7,119,149,1 61,190,238	Anti-fungal activity

22.	Isocalamendiol 2	10.85	C <sub>15</sub> H <sub>26</sub> O <sub>2</sub>	238	238.193 28		55,81,111,1 59,202,223, 238	Anti-fertility activity and anti-bacterial activity
23.	Cinnami acid , 4-hydroxy -3- methoxy-{5- hydroxy-2- hydroxymethyl	12.38	C <sub>31</sub> H <sub>40</sub> O <sub>15</sub>	652	652.236 72		55,77,91,10 5,121,151,1 68,194,215, 296,330,35 4,386,418	Anti-inflammatory properties
24.	(-)Spathulenol -	13.57	C <sub>15</sub> H <sub>24</sub> O 0	220	220.182 715		55,69,91,11 9,159,187,2 05,220	Antioxidant and anti- inflammatory activities
25.	1- Heptatriacotan ol	13.92	C <sub>37</sub> H <sub>76</sub> O 4	536	536.589 62		55,81,95,14 7,161,190,2 29,244,257	Antioxidant, anticancer, anti inflammatory and sex hormone activity
26.	Desacetylanqui dine	14.47	C <sub>17</sub> H <sub>24</sub> O <sub>6</sub> 4	324	324.157 288		55,67,91,10 5,124,145,1 59,187,205, 264,281	New chemical compound
27.	5-Isopropyl- 2,8-dimethyl- 9- oxatricyclo[4.4. .0.0.(2,8)]deca n-7-one	15.24	C <sub>14</sub> H <sub>23</sub> O <sub>2</sub> 0	222	222.161 98		55,69,81,95 ,109,123,15 1,162,194,2 22	Anti-termitic Activity
28.	Estra- 1,3,5(10)-trien- 17 $\beta$ -ol	15.81	C <sub>18</sub> H <sub>24</sub> O 8	256	256.182 714		57,73,85,97 ,129,157,18 5,213,241,2 56	Anti- osteoporosis activity
29.	Trans-1,2- Diaminocyclo hexane- N,N,N',N'- tetraacetic acid,	16.25	C <sub>14</sub> H <sub>22</sub> N <sub>2</sub> O <sub>8</sub> 3	346	346.137 615		55,81,96,11 0,153,181,1 97,225,241, 270	Unknown
30.	Phytol	16.66	C <sub>20</sub> H <sub>40</sub> O 5	296	296.307 917		57,71,81,95 ,111,123,13 7,196,221,2 49,278	Antioxidant activity
31.	Piperidine,1- (1-oxo-3- phenyl-2- propenyl)-	16.88	C <sub>14</sub> H <sub>17</sub> NO 3	215	215.131 014		51,84,103,1 31,149,172, 198,215	Antibacterial, analgesic and anti- inflammatory activity
32.	Eicosanoic acid , (acetoxy0-1- [(acetoxy)me thyl]ethyl ester	18.14	C <sub>27</sub> H <sub>50</sub> O <sub>6</sub> 2-7	470	470.360 739		57,71,84,98 ,117,137,15 9,171,227,2 69,295	Anti- inflammatory, and anti- melasma properties

33.	2,5,5,8a-Tetramethyl-6,7,8a-tetrahydro-5H-chromen-8-ol	18.53	C <sub>13</sub> H <sub>20</sub> O <sub>2</sub>	208	208.146 33		57,91,106,1 34,175,190, 208	Anticancer and antioxidant activity
34.	Z-5-methyl-6-heneicosen-11-one	18.55	C <sub>22</sub> H <sub>42</sub> O	322	322.323 566		55,71,83,97 ,139,169,19 7,241,307,3 22	Anti-Mycobacterium tuberculosis Activity, anti-inflammatory, analgesic and antimicrobial activities
35.	2H-1,2-Benzoxazine-3-carbonitrile,2-cyclohexyloctahydro-4a,8a-d	18.74	C <sub>17</sub> H <sub>28</sub> N <sub>2</sub> O	276	276.220 163		55,67,83,96 ,108,134,15 0,165,178,1 91,206,216, 234,249,27 6	Anti-inflammatory
36.	Indoxazin-4-one,4,5,6,7-tetrahydro-3-undecyl-	18.94	C <sub>18</sub> H <sub>29</sub> NO <sub>2</sub>	291	291.219 83		55,69,83,12 3,138,151,1 64,178,206, 234,262,29 1	New chemical compound
37.	9,10-Secocolestane-5,7,10(19)-triene-3,24,25-triol,(3 $\beta$ ,5 $Z$ ,7 $E$ )-	19.24	C <sub>27</sub> H <sub>44</sub> O <sub>3</sub>	416	416.329 044		55,118,136, 158,176,20 7,253,383,4 16	Biocides and anti-corrosion agents
38.	3-Oxo-10(14)-epoxyguai-11(13)-en-6,12-olide	19.37	C <sub>15</sub> H <sub>18</sub> O <sub>4</sub>	262	262.120 508		53,81,91,10 5,135,174,1 93,244,262	Anti-fungal activity
39.	9-Octadecenamide, n-butyl-	20.75	C <sub>22</sub> H <sub>43</sub> NO	337	337.334 465		55,69,100,1 28,184,210, 252,280,29 4,308,337 265,280,29 7,311,326	Antimicrobial, anti-inflammatory and antioxidant
40.	7-[2-(Ethoxycarbonyl)-3 $\alpha$ ,5 $\beta$ -dimethoxycyclopentyl]-1-heptanoic acid	20.96	C <sub>19</sub> H <sub>34</sub> O <sub>6</sub>	358	358.235 538		68,77,91,10 5,135,150,1 83,202,236, 264	New chemical compound
41.	2H-Benzo[f]oxiren-3-o[2,3-E]benzofuran-8,(9H)-one,9-[[1,3-benzodio-	21.24	C <sub>23</sub> H <sub>29</sub> NO <sub>5</sub>	399	399.204 573		81,91,115,1 28,152,174, 188,200,22 6,241,311	Antimicrobial and antifungal activities
42.	Nalorphine	21.27	C <sub>19</sub> H <sub>21</sub> NO <sub>3</sub>	311	311.152 143			Anti-pruritic agent against morphine

43.	2-Cyclohexen - 3-ol-1-one , 2- [1-iminotetradecyl],	21.73	C <sub>20</sub> H <sub>35</sub> NO <sub>2</sub>	321	321.266 779		55,83,110,1 25,153,166, 180,194,20 8,236,278,3 04,321 Antiulcer, hypolipidemic, antiatheroscler otic and anti- HIV
44.	Piperine	22.31	C <sub>17</sub> H <sub>19</sub> NO <sub>3</sub>	285	285.136 494		63,84,115,1 43,159,173, 201,229,25 Anti-pyretic and analgesic activities 6,285
45.	Fenretinide	22.78	C <sub>26</sub> H <sub>33</sub> NO <sub>2</sub>	391	391.251 13		58,69,81,95 109,135,14 8,161,202,2 Anti-proliferative activity 13,255,268
46.	11-Dehydrocorticosterone	24.67	C <sub>21</sub> H <sub>28</sub> O <sub>4</sub>	344	344.198 76		55,67,79,91 105,121,14 7,189,227,2 67,285,313, 344 Anti-proliferative activity
47.	5H-Cyclopropa[3,4]benz[1,2-e]azulen-5-one,1,1a,1b,4,4a,7a,7b	24.83	C <sub>20</sub> H <sub>28</sub> O <sub>5</sub>	348	348.193 674		55,83,173,1 99,217,312, 330,348 Biocides, anti- corrosion agents, drying agents, and coating
48.	17a-Ethyl-3 $\beta$ -methoxy-17a-aza-D-homoandrostan-5-ene-17-one	24.91	C <sub>22</sub> H <sub>35</sub> NO <sub>2</sub>	345	345.266 779		55,71,91,10 5,138,193,2 09,241,296, 330,345 Anti-cancer activity
49.	Bufa-20,22-dienolide m,14,15-epoxy-3,11-dihydroxy-(3 $\beta$ ,5 $\beta$ ,11 $\alpha$ ,15	25.11	C <sub>42</sub> H <sub>32</sub> O <sub>5</sub>	400	400.224 974		55,67,79,91 107,135,18 7,213,231,2 49,278,349, 364,382,40 0 Cell growth inhibitory activity, and antitumor
50.	9-Desoxo-9-x-acetoxy-3,8,12-tri-O-acetyltingol	25.52	C <sub>28</sub> H <sub>40</sub> O <sub>10</sub>	536	536.262 146		55,69,122,2 07,236,297, 357,417,47 7 Biocide and anti-fungal
51.	Retinal ,9-cis-3	25.96	C <sub>20</sub> H <sub>28</sub> O	284	284.214 016		55,69,79,95 119,133,17 3,199,213,2 41,255 Anti-oxidant role
52.	6- $\beta$ -Naltrexol	27.19	C <sub>20</sub> H <sub>25</sub> NO <sub>4</sub>	343	343.178 358		55,84,98,11 0,128,161,2 02,228,246, 288,302,32 4,343 Potential anti- angiogenic activity
53.	Ursodeoxycholic acid	28.34	C <sub>24</sub> H <sub>40</sub> O <sub>4</sub>	392	392.292 66		55,67,81,93 145,213,25 5,302,356,3 74,392 Antiinflammatory effect and anti- inflammatory activity



54.	5 $\alpha$ -Cholan-24-oic acid , 12 $\alpha$ -hydroxy-3,7-dioxo-,methyl ester	28.51	C <sub>25</sub> H <sub>38</sub> O <sub>5</sub>	418 925	418.271 925		55,95,219,245,285,345,382,400,418	Having biological activity (insect moulting hormones)
55.	Stigmasterol	29.16	C <sub>29</sub> H <sub>48</sub> O <sub>8</sub>	412 516	412.370 516		69,83,133,213,255,300,351,369,412	Anti-inflammatory and anti-parasitic effects

Table 2: FT-IR peak values of methanolic seeds extract of *Piper nigrum*.

S. No.	Peak (Wave number cm <sup>-1</sup> )	Intensity	Bond	Functional group assignment	Group frequency
1.	650.1	57.535	-	Unknown	-
2.	688.59	59.347	C-H	Alkenes	675-995
3.	704.02	60.135	C-H	Alkenes	675-995
4.	759.95	64.754	C-H	Alkenes	675-995
5.	827.46	67.689	C-H	Alkenes	675-995
6.	852.54	65.098	C-H	Alkenes	675-995
7.	894.97	68.412	C-H	Alkenes	675-995
8.	927.76	60.301	C-H	Alkenes	675-995
9.	999.13	38.637	-	Unknown	-
10.	1014.56	38.696	C-F stretch	Aliphatic fluoro compounds	1000-10150
11.	1076.28	54.627	C-O	Alcohols, Ethers, Carboxlic acids, Esters	1050-1300
12.	1134.14	61.967	C-O	Alcohols, Ethers, Carboxlic acids, Esters	1050-1300
13.	1149.57	63.780	C-O	Alcohols, Ethers, Carboxlic acids, Esters	1050-1300
14.	1195.87	71.015	C-O	Alcohols, Ethers, Carboxlic acids, Esters	1050-1300
15.	1249.87	55.761	C-O	Alcohols, Ethers, Carboxlic acids, Esters	1050-1300
16.	1292.87	72.884	C-O	Alcohols, Ethers, Carboxlic acids, Esters	1050-1300
17.	1361.74	68.851	NO <sub>2</sub>	Nitro Compounds	1300-1370
18.	1417.68	64.723	C-H	Alkanes	1340-1470
19.	1436.97	59.046	C-H	Alkanes	1340-1470
20.	1489.05	65.725	-	Unknown	-
21.	1506.41	66.871	-	Unknown	-
22.	1593.20	66.461	-	Unknown	-
23.	1616.35	64.515	-	Unknown	-
24.	1635.64	61.408	-	Unknown	-
25.	2665.62	88.750	-	Unknown	-
26.	2854.65	79.246	C-H	Alkanes	2850-2970
27.	2926.01	75.034	C-H	Alkanes	2850-2970

Table 3: Zone of inhibition (mm) of test bacterial strains to *Piper nigrum* bioactive compounds and standard antibiotics.  
/ *Piper nigrum* Antibiotics

	Bacteria				
	<i>Staphylococcus aureus</i>	<i>Escherichia coli</i>	<i>Proteus mirabilis</i>	<i>Klebsiella pneumonia</i>	<i>Pseudomonas eurogenosa</i>
<i>Piper nigrum</i>	4.00 $\pm$ 0.31	4.90 $\pm$ 0.13	5.00 $\pm$ 0.16	4.63 $\pm$ 0.41	4.12 $\pm$ 0.11
Rifambin	1.01 $\pm$ 0.10	0.77 $\pm$ 0.41	0.98 $\pm$ 0.11	1.00 $\pm$ 0.30	1.05 $\pm$ 0.42
Streptomycin	0.91 $\pm$ 0.27	1.60 $\pm$ 0.29	1.90 $\pm$ 0.10	0.96 $\pm$ 0.47	0.87 $\pm$ 0.20
Kanamycin	0.42 $\pm$ 0.18	1.12 $\pm$ 0.46	0.40 $\pm$ 0.12	1.00 $\pm$ 0.10	0.90 $\pm$ 0.47
Cefotoxime	0.87 $\pm$ 0.95	0.96 $\pm$ 0.27	0.93 $\pm$ 0.25	0.92 $\pm$ 0.18	0.71 $\pm$ 0.13

subsequent adverse effects<sup>6-12</sup>. Recently, biochemical activities of some important medicinal plants including *Piper* species and their metabolites have been described<sup>13-20</sup>. Pharmacological and clinical studies have revealed that piperine has CNS depressant, antipyretic, analgesic, anti-inflammatory (Ratner et al., 1991), antioxidant, and

hepatoprotective activities<sup>21,22</sup>. Studies have revealed anticonvulsant and bioavailability-enhancing properties of the drug. The fruits contain 1.0–2.5% volatile oil, 5–9% alkaloids, of which the major ones are piperine, chavicine, piperidine, and piperetine, and a resin<sup>23</sup>. The present study involves an assessment using GC-MS and FT-IR




Figure 1: GC-MS chromatogram of methanolic extract of *Piper nigrum*.

spectroscopic techniques to investigate the chemical composition methanolic fruit extract of *P. nigrum*.

## MATERIALS AND METHODS

### *Plant Material and Preparation of Extracts*

*P. nigrum* dried fruits were purchased from local market in Hilla city, middle of Iraq. after thorough cleaning and removal foreign materials, the fruits were stored in airtight container to avoid the effect of humidity and then stored at room temperature until further use. About 30 g of the plant sample powdered were soaked in 100 mL of methanol for 16 h in a rotatory shaker<sup>24,25</sup>. Whatman No.1 filter paper was used to separate the extract of plant. The filtrates were used for further phytochemical analysis. It was again filtered through sodium sulphate in order to remove the traces of moisture.

### *Identification of component by gas chromatography – mass spectrum analysis*

The physicochemical properties of the essential oil of *P. nigrum* L. (Black Pepper) are presented in Table1. Interpretation of mass spectroscopy (GC-MS) was conducted using data base of the National Institute Standard and Technology (NIST) having more than 62000 patterns. The spectrum of the unknown component was compared with the spectrum of the known component stored in the NIST library. The identity of the components in the extracts was assigned by the comparison of their retention indices and mass spectra fragmentation patterns with those stored on the computer library and also with published literatures<sup>26,27</sup>. The GC-MS analysis of the plant extract was made in a (Agilent 789 A) instrument under computer control at 70 eV. About 1µL of the methanol extract was injected into the GC-MS using a micro syringe and the scanning was done for 45 min. As the compounds were separated, they eluted from the column and entered a detector which was capable of creating an electronic signal whenever a compound was detected. The greater the concentration in the sample, bigger was the signal obtained which was then processed by a computer. The time from

when the injection was made (initial time) to when elution occurred is referred to as the retention time (RT). While the instrument was run, the computer generated a graph from the signal called chromatogram. Each of the peaks in the chromatogram represented the signal created when a compound eluted from the gas chromatography column into the detector. The x-axis showed the RT and the y-axis measured the intensity of the signal to quantify the component in the sample injected. As individual compounds eluted from the gas chromatographic column, they entered the electron ionization (mass spectroscopy) detector, where they were bombarded with a stream of electrons causing them to break apart into fragments. The fragments obtained were actually charged ions with a certain mass<sup>28</sup>. The M/Z (mass / charge) ratio obtained was calibrated from the graph obtained, which was called as the Mass spectrum graph which is the fingerprint of a molecule. Before analyzing the extract using gas chromatography and mass spectroscopy, the temperature of the oven, the flow rate of the gas used and the electron gun were programmed initially. The temperature of the oven was maintained at 100°C. Helium gas was used as a carrier as well as an eluent. The flow rate of helium was set to 1mL per min<sup>29-33</sup>. The electron gun of mass detector liberated electrons having energy of about 70eV. The column employed here for the separation of components was Elite 1(100% dimethyl poly siloxane).

### *Fourier transform infrared spectrophotometer (FTIR)*

The powdered sample of *P. nigrum* specimen was treated for FTIR spectroscopy (Shimadzu, IR Affinity 1, Japan). The sample was run at infrared region between 400 nm and 4000 nm<sup>31</sup>.

### *Determination of antibacterial activity of crude bioactive compounds of *Piper nigrum*.*

The test pathogens (*Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *E. coli*, and *Staphylococcus aureus*) were swabbed in Müller-Hinton agar plates. 60 µL of plant extract was loaded on the bored wells. The wells were




Figure 2: Structure of Cyclohexene, 1-methyl-5-(1-methylethenyl)-,(R) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 3: Structure of 1,6-Octadien-3-ol,3,7-dimethyl present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 4: Structure of 2-Methyl-1-ethylpyrrolidine present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 5: Structure of Propanedioic acid , dimethyl ester present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 6: Structure of Bicyclo[3.1.1]heptane, 6,6-dimethyl-2-methylene-,(1S) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 7: Structure of 3-Carene present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 8: Structure of Pyrrolizin-1,7-dione-6-carboxylic acid, methyl(ester) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 9: Structure of 7-epi-cis-sesquibabinene hydrate present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 10: Structure of Phenol, 2-methoxy-4-(1-propenyl)-, (Z) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 11: Structure of 2-Isopropenyl-5-methylhex-4-enal present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 12: Structure of L- $\alpha$ -Terpineol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 13: Structure of (R)-lavandulyl acetate present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 14: Structure of Epiglobulol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 15: Structure of Caryophyllene present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 16: Structure of 1,4,7-Cycloundecatriene, 1,5,9,9-tetramethyl-, Z,Z,Z- present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 17: Structure of Eugenol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 18: Structure of Alfa.Copaene present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 19: Structure of Naphthalene,1,2,3,5,6,7,8,8a-octahydro-1,8a-dimethyl-7-(1-methyl present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 20: Structure of Isocalamendiol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.





Figure 22: Structure of (-)-Spathulenol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.






Figure 24: Structure of  $\beta$ -copaene present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.






Figure 26: Structure of Estra-1,3,5(10)-trien-17 $\beta$ -ol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 27: Structure of Trans-1,2-Diaminocyclohexane-N,N,N',N'-tetraacetic acid present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 28: Structure of Phytol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 29: Structure of 1-Heptatriacanol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 30: Structure of Desacetylvanquidine present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 31: Structure of 5-Isopropyl-2,8-dimethyl-9-oxatricyclo[4.4.0.0(2,8)]decan-7-one present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 32: Structure of Z-5-methyl-6-heneicosen-11-one present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 33: Structure of 2H-1,2-Benzoxazine-3-carbonitrile, 2-cyclohexyloctahydro-4a,8a-d present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 34: Structure of Indoxazin-4-one, 4,5,6,7-tetrahydro-3-undecyl present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 35: Structure of Piperidine, 1-(1-oxo-3-phenyl-2-propenyl) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 36: Structure of Eicosanoic acid, 2-(acetoxy)-1-[(acetoxy)methyl]ethyl ester present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 37: Structure of 2,5,5,8a-Tetramethyl-6,7,8,8a-tetrahydro-5H-chromen-8-ol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 38: Structure of 7-[2-(Ethoxycarbonyl)-3 $\alpha$ ,5 $\beta$ -dimethoxycyclopentyl-1]-heptanoic acid present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 39: Structure of 2H-Benzofloxireno[2,3-E]benzofuran-8(9H)-one, 9-[[1,3-benzodio] present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 40: Structure of Nalorphine present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 41: Structure of 9,10-Secocoleta-5,7,10(19)-triene-3,24,25-triol, (3 $\beta$ ,5Z,7E) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 42: Structure of 3-Oxo-10(14)-epoxyguai-11(13)-en-6,12-olide present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 43: Structure of 9-Octadecenamide, n-butyl present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 44: Structure of 11-Dehydrocorticosterone present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 45: Structure of 5H-Cyclopropa[3,4]benz[1,2-e]azulen-5-one,1,1a,1b,4,4a,7a,7b present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 46: Structure of 17a-Ethyl-3 $\beta$ -methoxy-17a-aza-D-homoandrost-5-ene-17-one present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 47: Structure of 2-Cyclohexen-3-ol-1-one , 2-[1-iminotetradecyl] present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 48: Structure of Piperine present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 49: Structure of Fenretinide present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 50: Structure of 6- $\beta$ -Naltrexol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 51: Structure of Piperine present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 52: Structure of Ursodeoxycholic acid present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 53: Structure of Bufa-20,22-dienolide, 14,15-epoxy-3,11-dihydroxy-(3 $\beta$ ,5 $\beta$ ,11 $\alpha$ ,15 $\beta$ ) present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 54: Structure of 9-Desoxo-9-x-acetoxy-3,8,12-tri-O-acetyltingol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 55: Structure of Retinal, 9-cis present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 56: Structure of Stigmasterol present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.




Figure 57: Structure of 5 $\alpha$ -Cholan-24-oic acid, 12 $\alpha$ -hydroxy-3,7-dioxo-, methyl ester present in the methanolic seeds extract of *Piper nigrum* by using GC-MS analysis.



Figure 58: FT-IR profile of *Piper nigrum*.

bored in 0.5cm in diameter. The plates were incubated at 37°C for 24 h and examined. After the incubation the diameter of inhibition zones around the discs was measured.

## RESULTS AND DISCUSSION

Gas chromatography and mass spectroscopy analysis of compounds was carried out in methanolic fruits extract of *P. nigrum*, shown in Table 1. The GC-MS chromatogram of the 55 peaks of the compounds detected was shown in Figure 1. Chromatogram GC-MS analysis of the methanol extract of *P. nigrum* showed the presence of fifty five major peaks and the components corresponding to the peaks were determined as follows. The first set up peak were determined to be  $\alpha$ -pinene Figure 2. The second peak indicated to be Camphene Figure 3. The next peaks considered to be Eucalyptol, 2-Methoxy-4-vinylohenol, 1-Oxaspiron [4,5] deca-3,6-diene, 2,6,10,10-tetramethyl, 1-Oxaspiron [4,5]deca-3,6-diene, 2,6,10,10-tetramethyl, Neocurdione, Isoaromadendrene epoxide, 1b,4a-Epoxy-2H-cyclopenta[3,4]cyclopropa[8,9] cycloundec., Cis-

Vaccenic acid, Phenanthrenol, 4b,5,6,7,8,8a,9,10-octahydro-4b,8,8-trimethyl-1, Galanthamine, Dibenz[a,c]cyclohexane, 2,4,7-trimethoxy, 2,4a,7-Trihydroxy-1-methyl-8-methyleneqibb-3-ene, 1,10-carboxylic acid, Retinoic acid, 7,8,12-Tri-O-acetyl-3-desoxy-ingol-3-one, 4,6-Androstadien-3 $\beta$ -ol-17-one, acetate. (Figure 4-57). FTIR analysis of dry methanolic extract of *Rosmarinus officinalis* leaves proved the presence of Alkenes, Aliphatic fluoro compounds, Alcohols, Ethers, Carboxic acids, Esters and Nitro Compounds which shows major peaks at 688.59, 827.46, 927.76, 1014.56, 1249.87, 1361.74, 1417.68, 1506.41, 2665.62, 2854.65 and 2926.01 (Table 2; Figure 58). Boutekedjiret et al. (2003)<sup>32</sup> studied the constituents of rosemary essential oil from Algeria. They reported 1, 8-cineole, camphor,  $\beta$ -pinene, and  $\alpha$ -Pinene as the major constituents in the oil. Viuda-Martos et al. (2007)<sup>33</sup> investigated chemical composition of the essential oil of anther sample of rosemary leaves from Spain. The major constituents identified were  $\alpha$ -pinene, camphor, 1,8-cineole and camphene. The main components detected in

the oils were:  $\alpha$ -pinene, 1, 8-cineole, camphene, camphor, myrcene and broneol<sup>34</sup>. GC and GC-MS analysis of oils from rosemary leave samples from India revealed the presence of camphor, 1,8-cineole and  $\alpha$ -pinene as major constituents in the oils<sup>35-37</sup>. In this study five clinical pathogens selected for antibacterial activity namely, (*staphylococcus aeureus*, *klebsiella pneumoniae*, *pseudomonas aeruginosa*, *E. coli*. and *Proteus mirabilis*. Maximum zone formation against *Proteus mirabilis*, Table 3.

## CONCLUSION

*P. nigrum* is native plant of Iraq. It contain chemical constitutions which may be useful for various herbal formulation as anti-inflammatory, analgesic, antipyretic, cardiac tonic and antiasthamatic.

## ACKNOWLEDGEMENT

I thank Dr. Abdul-Kareem Al-Bermani, Lecturer, Department of Biology, for valuable suggestions and encouragement.

## REFERENCES

- Srinivas VP, Ashok KT, Uma MS, Anuradha V, Hari B. HPLC assisted chemo biological standardization of  $\alpha$ -glucosidase-Ienzyme inhibitory constituents from *Piper longum* Linn-An Indian medicinal plant, Journal of Ethno pharmacology. 2006; 108: 445–449.
- Nalini N, Manju V, Menon VP. Effect of spices on lipid metabolism in 1,2-dimethylhydrazine-induced rat colon carcinogenesis, J Med Food. 2006; 9(2): 237- 45.
- Altameme HJ, Hameed IH, Abu-Serag NA. Analysis of bioactive phytochemical compounds of two medicinal plants, *Equisetum arvense* and *Alchemila vulgaris* seed using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. Malays. Appl. Biol. 2015b; 44(4): 47– 58.
- Hamza LF, Kamal SA, Hameed IH. Determination of metabolites products by *Penicillium expansum* and evaluating antimicrobial activity. Journal of Pharmacognosy and Phytotherapy. 2015; 7(9): 194– 220.
- Hameed IH, Altameme HJ, Idan SA. Artemisia annua: Biochemical products analysis of methanolic aerial parts extract and anti-microbial capacity. Research Journal of Pharmaceutical, Biological and Chemical Sciences. 2016; 7(2): 1843- 1868.
- Catella-Lawson F, Reilly MP, Kapoor SC, Cucchiara AJ, DeMarco S, Tournier B, Vyas SN, FitzGerald GA. Cyclooxygenase inhibitors and the antiplatelet effects of aspirin. N. Engl. J. Med. 2001; 345: 1809–1817.
- Kawano KI, Hokamura K, Kondo K, Ikeda Y, Suzuki Y, Umemura K. Thromboxane A2 synthase inhibitor enhanced antithrombotic efficacy of GPIIb– IIIa receptor antagonist without increasing bleeding. Eur. J.Pharmacol. 2001; 417: 217–222.
- Hameed IH, Hamza LF, Kamal SA. Analysis of bioactive chemical compounds of *Aspergillus niger* by using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. Journal of Pharmacognosy and Phytotherapy. 2015b; 7(8): 132– 163.
- Mitsuhashi M, Tanaka A, Fujisawa C, Kawamoto K, Itakura A. Necessity of thromboxane A2 for initiation of platelet-mediated contact sensitivity: dual activation of platelets and vascular endothelial cells. J. Immunol. 2001; 166: 617–623.
- Dorsam RT, Kim S, Murugappan S, Rachoor S, Shankar H, Jin J, Kunapuli SP. Differential requirements for calcium and Src family kinases in platelet GPIIb/IIIa activation and thromboxane generation downstream of different G-protein pathways. Blood. 2005; 105: 2749–2756.
- Huang JS, Ramamurthy SK, Lin X, Le Breton GC. Cell signaling through thromboxane A2 receptors. Cell Signal. 2009; 16: 521–533.
- Hameed IH, Hussein HJ, Kareem MA, Hamad NS. Identification of five newly described bioactive chemical compounds in methanolic extract of *Mentha viridis* by using gas chromatography-mass spectrometry (GC-MS). Journal of Pharmacognosy and Phytotherapy. 2015c; 7 (7): 107-125.
- Nongyao S, Kitja S, Pathana P. Effects of *Piper longum* fruit, *Piper sarmentosum* root and *Quercus infectoria* nut gall on caecal amoebiasis in mice. Journal of Ethno pharmacology. 2004; 91: 357–360.
- Kartick M, Stanely MP. Preventive effect of rutin,a bioflavonoid, on lipid peroxides and antioxidants in isoproterenol-induced myocardial infarction in rats. J.Pharmacol. 2006; 58: 701-707.
- Altameme H J, Hadi MY, Hameed IH. Phytochemical analysis of *Urtica dioica* leaves by fourier-transform infrared spectroscopy and gas chromatography-mass spectrometry. Journal of Pharmacognosy and Phytotherapy. 2015a; 7(10): 238-252.
- Masaya I, Nobuaki O, Satoko, O, Masaki S, Norimichi N. Piperlongumine, a constituent of *Piper longum* L., inhibits rabbit platelet aggregation as a thromboxane A2 receptor antagonist, European Journal of Pharmacology. 2007; 570: 38-42.
- Seung WL, Young KK, Koanhoi K, Hyun SL, Jung HC, Woo SL. Alkamides from the fruits of *Piper longum* and *Piper nigrum* displaying potent cell adhesion inhibition. Bioorganic & Medicinal Chemistry Letters. 2008; 18: 4544-4546.
- Hameed IH, Ibraheem IA, Kadhim HJ. Gas chromatography mass spectrum and fourier-transform infrared spectroscopy analysis of methanolic extract of *Rosmarinus officinalis* leaves. Journal of Pharmacognosy and Phytotherapy. 2015d; 7 (6): 90- 106.
- Jagdale SC, Kuchekar BS, Chabukkar AR, Dlokhande P, Raut CG. Anti-Oxidant activity of *piper longum* L. Int.J.Biological chemistry. 2009; 3(3): 119-125.
- Al-Marzoqi AH, Hadi MY, Hameed IH. Determination of metabolites products by *Cassia angustifolia* and evaluate antimicrobial activity. Journal of Pharmacognosy and Phytotherapy. 2016; 8(2): 25-48.

21. Al-Marzoqi AH, Hameed IH, Idan SA. Analysis of bioactive chemical components of two medicinal plants (*Coriandrum sativum* and *Melia azedarach*) leaves using gas chromatography-mass spectrometry (GC-MS). African Journal of Biotechnology. 2015; 14(40): 2812-2830.
22. Hussein AO, Mohammed GJ, Hadi MY, Hameed IH. Phytochemical screening of methanolic dried galls extract of *Quercus infectoria* using gas chromatography-mass spectrometry (GC-MS) and Fourier transform-infrared (FT-IR). Journal of Pharmacognosy and Phytotherapy. 2016; 8(3): 49-59.
23. Park I, Lee S, Shin S, Park J, Young-Joon A. Larvicidal activity of isobutyl amides in *Piper nigrum* fruits against three mosquito species. Journal of Agriculture and Food Chemistry. 2002; 50: 1866–1870.
24. Altameme HJ, Hameed IH, Idan SA, Hadi MY. Biochemical analysis of *Origanum vulgare* seeds by fourier-transform infrared (FT-IR) spectroscopy and gas chromatography-mass spectrometry (GC-MS). Journal of Pharmacognosy and Phytotherapy. 2015c; 7(9): 221-237.
25. Jasim H, Hussein AO, Hameed IH, Kareem MA. Characterization of alkaloid constitution and evaluation of antimicrobial activity of *Solanum nigrum* using gas chromatography mass spectrometry (GC-MS). Journal of Pharmacognosy and Phytotherapy. 2015; 7 (4): 56-72.
26. Hussein HJ, Hadi MY, Hameed IH. Study of chemical composition of *Foeniculum vulgare* using Fourier transform infrared spectrophotometer and gas chromatography - mass spectrometry. Journal of Pharmacognosy and Phytotherapy. 2016; 8(3): 60-89.
27. Hadi MY, Mohammed GJ, Hameed IH. Analysis of bioactive chemical compounds of *Nigella sativa* using gas chromatography-mass spectrometry. Journal of Pharmacognosy and Phytotherapy. 2016; 8(2): 8-24.
28. Mohammed GJ, Al-Jassani MJ, Hameed IH. Antibacterial, Antifungal Activity and Chemical analysis of *Punica grantanum* (Pomegranate peel) using GC-MS and FTIR spectroscopy. International Journal of Pharmacognosy and Phytochemical Research. 2016; 8(3):480-494.
29. Hussein HM, Hameed IH, Ibraheem OA. Antimicrobial Activity and spectral chemical analysis of methanolic leaves extract of *Adiantum Capillus-Veneris* using GC-MS and FT-IR spectroscopy. International Journal of Pharmacognosy and Phytochemical Research. 2016; 8(3): 369-385.
30. Boutekedjiret C, Bentahar F, Belabbes R, Bessiere J. Extraction of rosemary essential oil by steam distillation and hydrodistillation. Flavour Fragr. J. 2003; 18: 481–484.
31. Viuda-Martos M, Yolanda R, Juana F, José P. Chemical Composition of the Essential Oils Obtained From Some Spices Widely Used in Mediterranean Region. Acta Chim. Slov. 2007; 54: 921–926.
32. Jamshidi R, Afzali Z, Afzali D. Chemical Composition of Hydrodistillation Essential Oil of Rosemary in Different Origins in Iran and Comparison with Other Countries. American-Eurasian J. Agric. & Environ. Sci. 2009; 5(1): 78-81.
33. Ram SV, Ur Rahman L, Sunita M, Rajesh KV Amit C, Anand S. Changes in essential oil content and composition of leaf and leaf powder of *Rosmarinus officinalis*. CIM-Hariyali during storage. Maejo International Journal of Science and Technology. 2011; 5(02): 181-19.
34. Kadhim MJ, Mohammed GJ, Hameed IH. In vitro antibacterial, antifungal and phytochemical analysis of methanolic fruit extract of *Cassia fistula*. Oriental Journal of Chemistry. 2016; 32(2): 10-30.
35. Shareef HK, Muhammed HJ, Hussein HM, Hameed IH. Antibacterial effect of ginger (*Zingiber officinale*) roscoe and bioactive chemical analysis using gas chromatography mass spectrum. Oriental Journal of Chemistry. 2016; 32(2): 20-40.
36. Al-Jassaci MJ, Mohammed GJ, Hameed IH. Secondary Metabolites Analysis of *Saccharomyces cerevisiae* and Evaluation of Antibacterial Activity. International Journal of Pharmaceutical and Clinical Research. 2016; 8(5): 304-315.
37. Aronow WS. Management of peripheral arterial disease of the lowerextremities in elderly patients. J. Gerontol. 2004; 59: 172–177.