

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/305930548>

Analysis of the Secondary Metabolite Products of Ammi majus and Evaluation Anti-Insect Activity

Article in *International Journal of Pharmacognosy and Phytochemical Research* · January 2016

READ

1

3 authors, including:



Imad Hadi Hameed

University of Babylon

61 PUBLICATIONS 325 CITATIONS

SEE PROFILE

Analysis of the Secondary Metabolite Products of *Ammi majus* and Evaluation Anti-Insect Activity

Haider Mashkooor Hussein¹, Imad Hadi Hameed^{2*}, Jenan Mohammed Ubaid³

¹College of Science, Al-Qadisiya University, Iraq.

²College of Nursing, Babylon University, Iraq.

³College of Science for women, Babylon University, Iraq.

Available Online: 10th August, 2016

ABSTRACT

Aims of our study were analysis of the secondary metabolite products of *Ammi majus* and evaluation insecticidal activity against *Tribolium castaneum*. The identification of bioactive chemical compounds is based on the peak area, retention time molecular weight and molecular formula. GC-MS analysis of *Ammi majus* revealed the existence of the Pregn-5-ene-3,11-dione, 17,20:20,21-bis [methylenebis(oxy)]-, Penta-2,4-dien-1-one, 5-dimethylamino-1-[5-(4-dimethylamino)], 8-Octadecenal, 9,10-Secocholesta-5,7,10(19)-triene-3,24,25-triol, (3 β ,5Z,7E)-, 3,7-Diacetamido-7H-s-triazolo[5,1-c]-s-triazole, Piperidine, 2,3-dimethyl-, d-Mannose, 1-Nitro-2-acetamido-1,2-dideoxy-d-mannitol, Paromomycin, d-Glycero-d-ido-heptose, Aminoacetamide, N-methyl-N-[4-(1-pyrrolidinyl)-2-butynyl]-, Pyrrolizin-1,7-dione-6-carboxylic acid, methyl(ester), 5-Allyl-6-methyl-2-phenyl-5,6-dihydro-4H-oxazolo[4,5-c]pyridin-, 3-O-Methyl-d-glucose, Dasycarpidan-1-methanol, acetate (ester) and Corynan-17-ol,18,19-didehydro-10-methoxy-,acetate (ester). Methanolic extract of *Ammi majus* was active on accumulative mortality of *Tribolium castaneum* (adult).

Keywords: *Ammi majus*, Anti-insect, Bioactive chemical analysis, GC/MS, *Tribolium castaneum*.

INTRODUCTION

Several plant derived medicine are rich in phenolic and other compounds such as those used in protection against the coronary heart diseases and carcinogenesis¹. Fructus *Ammi Majoris* consists of the dried ripe fruits of *Ammi majus* L. (Apiaceae)^{1,2}, and widely distributed in Europe, the Mediterranean region, and western Asia, now cultivated in India². *Ammi majus* has different names, the Arabic name khillah, khillah shyani, English name bishops weed, Latin and German name ammi, French name ammi commun, belongs to family Apiaceae. Is an upright annual herb which grows up to 1 meter or more, whitish tap-roots; stem erect, slender, leaves alternate with long petiole, flower whitish actinomorphic or zygomorphic, fruit a cremocarp with slender prominent ribs, *A. majus* is one of the best medicinal plants that was first found in Mediterranean and Egypt³. It is used as an emmenagogue to regulate menstruation, as a diuretic, and for treatment of leprosy, kidney stones, and urinary tract infections⁴. This is widely used for the treatment of skin disorders such as psoriasis and vitiligo (acquired leukoderma), and of vitiligo⁵. In Europe due to climate conditions the growth of *Ammi majus* is poor, attempts to acclimatize *Ammi majus* in cool climate were not successful, the fruits failed to ripen and plants were highly susceptible to infection⁶. The phenolic compounds such as flavonoids, phenolic acids and tannins are considered to be major contributors to the antioxidant capacity of plants⁷⁻⁹. The objectives of this study were

analysis of the secondary metabolite products and evaluation anti-insect activity against adults of *Tribolium castaneum*.

MATERIALS AND METHODS

Plant material, extraction and isolation of Ammi majus

One kg of *Ammi majus* have been collected from Hilla city, middle of Iraq. *Ammi majus* was stored in airtight container to avoid the effect of humidity and then stored at room temperature until further use. Methanolic extract of *Ammi majus* powdered were soaked in 500 mL methanol for 14 hours in a rotatory shaker. The filtrates were used for further phytochemical analysis¹⁰⁻¹⁶.

Evaluation of anti-insect activity









Tribolium castaneum was obtained from laboratory cultures maintained in the dark in incubators at 26 \pm 1°C. This insect was reared on wheat flour mixed with yeast (10:1, w:w). A control was prepared in the same way but extract application was omitted. Three replicates were set up for the treated. Results of the study were based on analysis of variance (ANOVA) using Statistica Software. A significance level of 0.05 was used for all statistical tests¹⁷⁻²².

Gas chromatography – mass spectrum analysis

The Gas chromatography – mass spectrum analysis of methanolic extract of *Ammi majus* was made in a (Agilent 789 A) instrument under computer control at 70 eV. About 1 μ L of the methanol extract was injected into the

Table 1: Major phytochemical compounds identified in methanolic extract of *Ammi majus*.

S. No	Phytochemical compound	RT (min)	Molecular Weight	Exact Mass	Chemical structure	MS Fragmentations	Pharmacological actions
1.	Pregn-5-ene-3,11-dione, 17,20:20,21-bis[methylenebis(oxy)]-	3.178	446	446.230453		55,69,81,99,161,256,314,372,446	Unknown
2.	Penta-2,4-dien-1-one, 5-dimethylamino-1-[5-(4-dimethylamino)]	3.402	302	302.145285		53,81,94,124,189,221,258,302	anti-inflammatory properties
3.	8-Octadecenal	3.556	266	266.260965		57,68,82,97,177,252	anti bacterial and anti fungal activities
4.	9,10-Secocholesta-5,7,10(19)-triene-3,24,25-triol (3β,5Z,7E)	3.653	416	416.329044		55,69,118,136,158,176,207,253,383,416	antitumor activity
5.	3,7-Diacetamido-7H-s-triazolo[5,1-c]-s-triazole	4.237	223	223.081773		59,69,84,97,153,207	anti-mycobacterium tuberculosis activity
6.	Piperidine, 2,3-dimethyl-	4.546	113	113.1204495		56,70,84,98	antibacterial and antifungal activity
7.	d-Mannose	4.666	180	180.063388		60,73,103,149	Anti-Bacterial Agents
8.	1-Nitro-2-acetamido-1,2-dideoxy-d-mannitol	4.832	252	252.095751		60,86,98,114,161,188,219,252	antifungal activity

9.	Paromomycin	5.055	615	615.296303		57,67,80,94,109,191,227,,259,292,324	<i>anti-HIV-1 agents</i>
10.	d-Glycero-d-ido-heptose	5.141	210	210.073953		60,73,85,133	<i>anti-inflammatory and anti-septic activities</i>
11.	Aminoacetamide, N-methyl-N-[4-(1-pyrrolidinyl)-2-butynyl]-	6.240	209	209.152812		55,70,84,121,139,192	Unknown
12.	Pyrrolizidin-1,7-dione-6-carboxylic acid, methyl(ester)	8.122	197	197.068808		55,69,84,98,142,197	Unknown
13.	5-Allyl-6-methyl-2-phenyl-5,6-dihydro-4H-oxazolo[4,5-c]pyridin-3-O-Methyl-d-glucose	7.739	268	268.121178		56,67,83,105,129,165,197,240	<i>anti-inflammatory, antitumor and anti-parasitic activity</i>
14.	3-O-Methyl-d-glucose	13.283	194	194.079039		73,87,103,116,145,163,177	<i>anti-cancer, anti-inflammatory</i>
15.	Dasycarpidan-1-methanol, acetate (ester)	16.877	326	326.199429		69,97,180,222,256,326	antimicrobial, antioxidant and <i>anti-inflammatory</i>
16.	Corynan-17-ol,18,19-didehydro-10-methoxy-,acetate (ester)	17.975	368	368.209993		69,97,143,186,281,308,368	<i>anti-diarrhoeal activity</i>

GC-MS using a micro syringe and the scanning was done for 45 minutes. The time from when the injection was made (Initial time) to when elution occurred referred to as the Retention time (RT)²³⁻³⁰. While the instrument was run, the computer generated a graph from the signal

called Chromatogram. Each of the peaks in the chromatogram represented the signal created when a compound eluted from the Gas chromatography column into the detector. The M/Z (mass / charge) ratio obtained was calibrated from the graph obtained, which was called



Figure 1: GC-MS chromatogram of methanolic extract of *Ammi majus*.

Figure 2: Mass spectrum of Pregnen-5-ene-3,11-dione , 17,20:20,21-bis[methylenebis(oxy)]- with Retention Time (RT)= 3.178.




Figure 3: Mass spectrum of Penta-2,4-dien-1-one , 5-dimethylamino-1-[5-(4-dimethylamino)] with Retention Time (RT)= 3.402.





Figure 4: Mass spectrum of 8-Octadecenal with Retention Time (RT)= 3.556.

Figure 5: Mass spectrum of 9,10-Secocholesta-5,7,10(19)-triene-3,24,25-triol , (3 β ,5Z,7E)- with Retention Time (RT)= 3.653.




Figure 6: Mass spectrum of 3,7-Diacetamido-7H-s-triazolo[5,1-c]-s-triazole with Retention Time (RT)= 4.237.




Figure 7: Mass spectrum of Piperidine, 2,3-dimethyl- with Retention Time (RT)= 4.546.




Figure 8: Mass spectrum of d-Mannose with Retention Time (RT)= 4.666.




Figure 9: Mass spectrum of 1-Nitro-2-acetamido-1,2-dideoxy-d-mannitol with Retention Time (RT)= 4.832.




Figure 10: Mass spectrum of Paromomycin with Retention Time (RT)= 5.055.




Figure 11: Mass spectrum of d-Glycero-d-ido-heptose with Retention Time (RT)= 5.141.




Figure 12: Mass spectrum of Aminoacetamide, N-methyl-N-[4-(1-pyrrolidinyl)-2-butynyl]- with Retention Time (RT)= 6.240.




Figure 13: Mass spectrum of Pyrrolizin-1,7-dione-6-carboxylic acid, methyl(ester) with Retention Time (RT)= 8.122.




Figure 14: Mass spectrum of 5-Allyl-6-methyl-2-phenyl-5,6-dihydro-4H-oxazolo[4,5-c]pyridin- with Retention Time (RT)= 7.739.




Figure 15: Mass spectrum of 3-O-Methyl-d-glucose with Retention Time (RT)= 13.283.




Figure 16: Mass spectrum of Dasycarpidan-1-methanol, acetate (ester) with Retention Time (RT)= 16.877.




Figure 17: Mass spectrum of Corynan-17-ol, 18,19-didehydro-10-methoxy-, acetate (ester) with Retention Time (RT)= 17.975.




Figure 18: Effect of methanolic leaves extract of *Ammi majus* on accumulative mortality of *Tribolium castaneum* (adult).

as the Mass spectrum graph which is the fingerprint of a molecule. Before analyzing the extract using gas chromatography and mass spectroscopy, the temperature of the oven, the flow rate of the gas used and the electron gun were programmed initially. The identity of the components in the extracts was assigned by the comparison of their retention indices and mass spectra fragmentation patterns with those stored on the computer library and also with published literatures. Compounds were identified by comparing their spectra to those of the Wiley and NIST/EPA/NIH mass spectral libraries³¹⁻³³.

RESULTS AND DISCUSSION

Characterization of phytochemical compounds

GC-MS is a powerful technique used for many applications which has very high sensitivity and specificity. GC-MS analysis of compounds was carried out in methanolic extract of *Ammi majus*, shown in Table 1. The GC-MS chromatogram of the 16 peaks of the compounds detected was shown in Figure 1. The First set up peak were determined to be Pregn-5-ene-3,11-dione, 17,20:20,21-bis [methylenebis(oxy)]-, Penta-2,4-dien-1-one, 5-dimethylamino-1-[5-(4-dimethylamino)], 8-Octadecenal, 9,10-Secocholesta-5,7,10(19)-triene-3,24,25-triol, (3 β ,5Z,7E)-, 3,7-Diacetamido-7H-s-triazolo[5,1-c]-s-triazole, Piperidine, 2,3-dimethyl-, d-Mannose, 1-Nitro-2-acetamido-1,2-dideoxy-d-mannitol, Paromomycin, d-Glycero-d-ido-heptose, Aminoacetamide, N-methyl-N-[4-(1-pyrrolidinyl)-2-butynyl]-, Pyrrolizin-1,7-dione-6-carboxylic acid, methyl(ester), 5-Allyl-6-methyl-2-phenyl-5,6-dihydro-4H-oxazolo[4,5-c]pyridin-, 3-O-Methyl-d-glucose, Dasycarpidan-1-methanol, acetate (ester) and Corynan-

17-ol,18,19-didehydro-10-methoxy-, acetate (ester) Figure 2-17.

Anti-insect activity

In the current study, the anti-insect activity of the methanolic extract was evaluated. Extract of *Ammi majus* caused 50% mortality during the 30 days after treatment. The methanol extracts of *Ammi majus* significantly affected survival of adult with 79%, during 45 days after treatment Figure 18. The relation between exposure period and treatment was very significant $p < 0.05$. Significant insecticidal activity against *T. castaneum* adults was observed with crude methanol extract from *Ammi majus*. Adults were more susceptible than larvae to extract of *Ammi majus*. Beside antimicrobial activity, the essential oil and its constituents have also been used for their herbicidal²⁶⁻³⁵, insecticidal, and anti-lice properties, as well as in integrated disease management against phytopathogenic fungi, nonspecific skin infections. *Ammi majus* belong to the family Apiaceae that provide a huge number of compounds with important medicinal activities. Numerous clinical trials have assessed the efficacy of fructus *Ammi Majoris* and xanthotoxin for the treatment of vitiligo, psoriasis, and hypopigmentation tinea versicolor. The natural source of producing huge numbers of phytoconstituents in a most efficient way and with precise selectivity are plants and different bioactive phytoconstituents have been isolated and characterized in the middle of the 19th century, many of these are used as the active ingredients of the modern medicine or as the lead compounds for new drug discovery³⁶⁻⁴³.

CONCLUSION

Sixteen bioactive chemical constituents have been identified from methanolic extract of the *Ammi majus* by

Gas Chromatography-Mass Spectrometry technique. Evaluation of anti-insect of secondary metabolite products of *Ammi majus* forms a primary platform for further pharmacological investigation for the development of new potential anti-insect compounds.

ACKNOWLEDGEMENT

The authors are gratefully acknowledging a head of department of biology, College of Science for women, Babylon University.

REFERENCES

1. El-Mofty AM, El-Sawalhy H, El-Mofty M. Clinical study of a new preparation of 8-methoxypsoralen in photochemotherapy. *Int J Dermatol*. 1994; 33:588–592.
2. Vijayalakshmi R, Ravindhran R. Preliminary comparative phytochemical screening of root extracts of *Diospyrus ferrea* (Wild.) Bakh and *Aerva lanata* (L.) Juss. Ex Schultes. *Pelagia Research Library Asian J of Plant Sci Res* 2012; 2(5): 581-587.
3. Krolicka A, Staniszewska I, Bielawski K, Malinski E, Szafraniek J, ojkowska E. Establishment of hairy root cultures of *Ammi majus*. *Plant Sci* 2001; 160: 259-264.
4. Khan NG, Khawas-ul-Advia, Khadim-ul-taleem, Steam Press, Lahore 1991; 1:190.
5. Lin J, Zhang SM, WuK, Willett WC, Fuchs CS, Giovannucci E. Flavonoid Intake and colorectal cancer in men and women. *Amer J of Epidem* 2006; 164:644-651.
6. Al-Hadidi AK, Al-Numan YA, Al-Daody CA. Interaction between some phenolic compounds in *ammi majus* herb (khillah) extracts and antibiotics against some selected bacterial isolates *in vitro*. *Raf J Sci*. 2013; 24(2):17-30.
7. El-Mofty AM, El-Sawalhy H, El-Mofty M. Photochemotherapy in the treatment of post tinea versicolor hypopigmentation. *Med J Cairo Univ*. 1995; 61(4):632–637.
8. Fakim GA. Medicinal plants: traditions of yesterday and drugs of tomorrow. *Mol Aspects Med*. 2006; 7(1):1-93.
9. Erkan N, Ayranci G, Ayranci E. Antioxidant activity of rosemary (*Rosmarinus officinalis*) extract, Black seed (*Nigella sativa*) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food Chem*. 2008; 110:76-82.
10. Kadhim MJ, Sosa AA, Hameed IH. Evaluation of anti-bacterial activity and bioactive chemical analysis of *Ocimum basilicum* using Fourier transform infrared (FT-IR) and gas chromatography-mass spectrometry (GC-MS) techniques. *International Journal of Pharmacognosy and Phytochemical Research*. 2016; 8(6): 127-146.
11. Mohammed GJ, Kadhim MJ, Hussein HM. Characterization of bioactive chemical compounds from *Aspergillus terreus* and evaluation of antibacterial and antifungal activity. *International Journal of Pharmacognosy and Phytochemical Research*. 2016; 8(6): 889-905.
12. Hameed IH, Altameme HJ, Idan SA. *Artemisia annua*: Biochemical products analysis of methanolic aerial parts extract and anti-microbial capacity. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2016; 7(2): 1843- 1868.
13. Hussein AO, Mohammed GJ, Hadi MY, Hameed IH. Phytochemical screening of methanolic dried galls extract of *Quercus infectoria* using gas chromatography-mass spectrometry (GC-MS) and Fourier transform-infrared (FT-IR). *Journal of Pharmacognosy and Phytotherapy*. 2016; 8(3): 49-59.
14. Sosa AA, Bagi SH, Hameed IH. Analysis of bioactive chemical compounds of *Euphorbia lathyris* using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. *International Journal of Pharmacognosy and Phytochemical Research*. 2016; 8(5): 109-126.
15. Altameme H J, Hadi MY, Hameed IH. Phytochemical analysis of *Urtica dioica* leaves by fourier-transform infrared spectroscopy and gas chromatography-mass spectrometry. *Journal of Pharmacognosy and Phytotherapy*. 2015a; 7(10): 238-252.
16. Lamberty M, D Zachary, R Lanot, Bordereau C, Robert A, Hoffmann JA, Bulet P. Insect immunity. Constitutive expression of a cysteine-rich antifungal and a linear antibacterial peptide in a termite insect. *Journal of Biology Chemistry*. 2001; 276: 4085-4092.
17. Shukla R, Siravatava B, Kumar R, Dubey NK. Potential of some powders in reducing infection of chickpea by *Callosobruchus maculatus* (Coleoptera: Brauchidae). *J. Agricultur. Technol*. 2007; 3(1): 11-19.
18. Mohammed GJ, Omran AM, Hussein HM. Antibacterial and Phytochemical Analysis of *Piper nigrum* using Gas Chromatography-Mass Spectrum and Fourier-Transform Infrared Spectroscopy. *International Journal of Pharmacognosy and Phytochemical Research*. 2016; 8(6): 977-996.
19. Hamza LF, Kamal SA, Hameed IH. Determination of metabolites products by *Penicillium expansum* and evaluating antimicrobial activity. *Journal of Pharmacognosy and Phytotherapy*. 2015; 7(9): 194-220.
20. Jasim H, Hussein AO, Hameed IH, Kareem MA. Characterization of alkaloid constitution and evaluation of antimicrobial activity of *Solanum nigrum* using gas chromatography mass spectrometry (GC-MS). *Journal of Pharmacognosy and Phytotherapy*. 2015; 7(4): 56-72.
21. Hadi MY, Mohammed GJ, Hameed IH. Analysis of bioactive chemical compounds of *Nigella sativa* using gas chromatography-mass spectrometry. *Journal of Pharmacognosy and Phytotherapy*. 2016; 8(2): 8-24.
22. Hameed IH, Ibraheam IA, Kadhim HJ. Gas chromatography mass spectrum and fourier-transform infrared spectroscopy analysis of methanolic extract of *Rosmarinus oficinalis* leaves. *Journal of*

- Pharmacognosy and Phytotherapy. 2015; 7 (6): 90-106.
23. Shareef HK, Muhammed HJ, Hussein HM, Hameed IH. Antibacterial effect of ginger (*Zingiber officinale*) roscoe and bioactive chemical analysis using gas chromatography mass spectrum. *Oriental Journal of Chemistry.* 2016; 32(2): 20-40.
24. Al-Jassaci MJ, Mohammed GJ, Hameed IH. Secondary Metabolites Analysis of *Saccharomyces cerevisiae* and Evaluation of Antibacterial Activity. *International Journal of Pharmaceutical and Clinical Research.* 2016; 8(5): 304-315.
25. Mohammed GJ, Al-Jassani MJ, Hameed IH. Antibacterial, Antifungal Activity and Chemical analysis of *Punica grantanum* (Pomegranate peel) using GC-MS and FTIR spectroscopy. *International Journal of Pharmacognosy and Phytochemical Research.* 2016; 8(3): 480-494.
26. Al-Marzoqi AH, Hadi MY, Hameed IH. Determination of metabolites products by *Cassia angustifolia* and evaluate antimicrobial activity. *Journal of Pharmacognosy and Phytotherapy.* 2016; 8(2): 25-48.
27. Altameme HJ, Hameed IH, Abu-Serag NA. Analysis of bioactive phytochemical compounds of two medicinal plants, *Equisetum arvense* and *Alchemilla vulgaris* seed using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. *Malays. Appl. Biol.* 2015b; 44(4): 47-58.
28. Hameed IH, Hamza LF, Kamal SA. Analysis of bioactive chemical compounds of *Aspergillus niger* by using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. *Journal of Pharmacognosy and Phytotherapy.* 2015; 7(8): 132-163.
29. Hameed IH, Hussein HJ, Kareem MA, Hamad NS. Identification of five newly described bioactive chemical compounds in methanolic extract of *Mentha viridis* by using gas chromatography-mass spectrometry (GC-MS). *Journal of Pharmacognosy and Phytotherapy.* 2015; 7 (7): 107-125.
30. Hussein HM, Hameed IH, Ibraheem OA. Antimicrobial Activity and spectral chemical analysis of methanolic leaves extract of *Adiantum Capillus-Veneris* using GC-MS and FT-IR spectroscopy. *International Journal of Pharmacognosy and Phytochemical Research.* 2016; 8(3): 369-385.
31. Hussein HJ, Hadi MY, Hameed IH. Study of chemical composition of *Foeniculum vulgare* using Fourier transform infrared spectrophotometer and gas chromatography - mass spectrometry. *Journal of Pharmacognosy and Phytotherapy.* 2016; 8(3): 60-89.
32. Kadhim MJ, Mohammed GJ, Hameed IH. In vitro antibacterial, antifungal and phytochemical analysis of methanolic fruit extract of *Cassia fistula*. *Oriental Journal of Chemistry.* 2016; 32(2): 10-30.
33. Altameme HJ, Hameed IH, Idan SA, Hadi MY. Biochemical analysis of *Origanum vulgare* seeds by fourier-transform infrared (FT-IR) spectroscopy and gas chromatography-mass spectrometry (GC-MS). *Journal of Pharmacognosy and Phytotherapy.* 2015c; 7(9): 221-237.
34. Yasuhara-Bell J, Yang Y, Barlow R, Trapido-Rosenthal H, Lu Y. In vitro evaluation of marine-microorganism extracts for anti-viral activity. *Virol.* 2010; 7:182.
35. Soltan MM, Zaki AK. Antimicrobial and antiviral activities of some Egyptian medicinal plants. *J Ethnopharmacol.* 2009; 126(1): 102-107.
36. Hussein HM. Determination of phytochemical composition and ten elements content (CD, CA, CR, CO, FE, PB, MG, MN, NI AND ZN) of *CARDARIA DRABA* by GC-MS, FT-IR and AAS technique. *Int. J Pharm Bio Sci.* 2016;7(3): (B) 1009 – 1017.
37. Hussein HM. Analysis of trace heavy metals and volatile chemical compounds of *Lepidium sativum* using atomic absorption spectroscopy, gas chromatography-mass spectrometric and fourier-transform infrared spectroscopy. *Research Journal of Pharmaceutical, Biological and Chemical Sciences.* 2016;7(4): 2529 – 2555.
38. Jaddoa HH, Hameed IH, Mohammed GJ. Analysis of volatile metabolites released by *Staphylococcus aureus* using gas chromatography-Mass spectrometry and determination of its antifungal activity. *Oriental Journal of Chemistry.* 2016;32(4).
39. Hameed IH, Salman HD, Mohammed GJ. Evaluation of antifungal and antibacterial activity and analysis of bioactive phytochemical compounds of *Cinnamomum zeylanicum* (Cinnamon bark) using gas chromatography-mass spectrometry. *Oriental Journal of Chemistry.* 2016;32(4).
40. Sahi NM. Evaluation of insecticidal activity of bioactive compounds from *Eucalyptus citriodora* against *Tribolium castaneum*. *International Journal of Pharmacognosy and Phytochemical Research.* 2016; 8(8).
41. Ubaid JM, Hussein HM, Hameed IH. Determination of bioactive chemical composition of *Callosobruchus maculatus* and investigation of its anti-fungal activity. *International Journal of Pharmacognosy and Phytochemical Research.* 2016; 8(8).
42. Kadhim MJ, Mohammed GJ, Hussein HM. Analysis of bioactive metabolites from *Candida albicans* using (GC-MS) and evaluation of antibacterial activity. *International Journal of Pharmaceutical and Clinical Research.* 2016; 8(7): 655-670.
43. Ubaid JM, Hussein HM, Hameed IH. Analysis of bioactive compounds of *Tribolium castaneum* and evaluation of anti-bacterial activity. *International Journal of Pharmaceutical and Clinical Research.* 2016; 8(7): 655-670.