

Prevalence β -Lactamase broad-spectrum type TEM between isolates of *Proteus mirabilis* in AL-Diwaniya city.

Nuha Jawad and Syoof Khowman

Biology Department, College of Sciences, AL- Qadisiya University, AL-Diwanyia province, Iraq.

abstract

The samples of the study were collected from different clinical sources 185 isolates of hospitals, the Diwaniyah city during the period from October 2015 to April and divided the samples, according to sources collected into five groups (46 swab ear, 42 swab burns, 37 a stool sample, 59 urine sample and 1 blood sample), as results showed that cultural and biochemical tests 69 isolated belong to bacteria *Proteus mirabilis*, diagnosis was confirmed by api 20E and the use of polymerase chain reaction. In the study showed that there are (60.86%) isolates were resistant these isolates vowed multi-resistant of antibiotics (multidrug resistance) the highest proportion among the three types of resistance, and the isolates of the overall resistance (extensive drug resistance) ratio of resistance (34.78 %) , while the third type of resistance the resistance ratio was increased by two (4.34 %) and represents the resistance of each species of antibiotics in the current study (pand drug resistance) . Tested 24 isolates *P.mirabilis* to examine the capability of production of enzymes β - lactam TEM type using PCR, was *bla*_{TEM} -177 the most frequently identified gene among isolates of *P.mirabilis* percentage (87.5%), while the *bla*_{TEM} -160 (59.38%), and *bla*_{TEM} -72 (56.25%), *bla*_{TEM} -1(56.25%), *bla*_{TEM} -156 (53.13%), *bla*_{TEM} -89 (31.25%), while *bla*_{TEM} -3 counted the lowest percentage among the studied genes in *P.mirabilis* ratio (6.25%). Analyzed Phylogenetic tree genetic tree analysis *bla*_{TEM} by Mega6 program, were used genetic tree analysis of the type UPGMATree Test yields 14 sample of isolates *P.mirabilis*. compared 2 isolates of *P.mirabilis* multi-resistant of antibiotics and producing resistance gene type TEM β -lactamase to analyze the genetic tree. The results of the analysis, there is a clear Identify of isolates of *P.mirabilis bla*_{TEM} local isolates with worldwide origin *P.mirabilis* compared with the rest of the other species that appear in the genetic tree analysis.

Introduction

P. mirabilis and one of the most important bacterial species negative bacteria belong to the family of intestinal and these bacteria of great significance in the spread of hospital-acquired infections Nosocomial infections (1),(2) and is one of the most common pathogens that cause infections acquired community community acquired infections, and for being opportunistic bacteria opportunistic therefore causing many injuries pathological when its presence is in their natural habitat (2),(3).The resistant of *P.mirabilis* especially multi-drug antibiotic resistance become developing with time as well as the production of this type of enzymes β -lactamase, most of the genes multiple drug resistance antibiotics is transmitted by plasmids (4). β -lactamase broad spectrum enzymes produced by negative bacteria It gives the bacteria produced an increase in resistance to antibiotics commonly used (5). So is the study of β -lactamase enzymes is very important and necessary to decrease the production and spread of β -lactamase(6). These enzymes cause failed treatments to control the various bacterial

infections, for resistance of antibiotics interested researchers (7) these enzymes become increasing gradually with time and is developing rapidly (8)

Materials and Methods:

Collection of Samples: During the period from October 2015 to April 2016, a total of 185 nonduplicate clinical samples in Al-Diwaniya city. Samples which were collected by sterile swabs and containers had been cultured on blood agar and macConkey agar, to get pure colonies subculture done on macConkey agar, incubated for overnight at 37C°.

Polymerase Chain Reaction Amplification (PCR): was used to screen for the occurrence the genes *bla_{TEM}* in *P.mirabilis* that resistance to antibiotic β -lactam : , *bla_{TEM}*-177, *bla_{TEM}*-89, *bla_{TEM}*-156 , *bla_{TEM}*-160 *bla_{TEM}*-72 , *bla_{TEM}*-3, *bla_{TEM}*-1 . the Primers were designed from sequences deposited in the GenBank database Table (1). The complete template DNA for the PCR amplification was extracted from the supernatant of a combination of *P.mirabilis* cells produced by salting out method (9). PCR amplification was using 5 μ l of the template DNA, 2 μ l of each primer, 10 μ l master mix, and 1 μ l of Taq DNA polymerase in a total volume of 20 μ l. A thermocycler (Mastercycler gradient; Eppendorf, Hamburg, Germany) was programmed with the suitable conditions (9). Then, 5 μ l of each PCR product was examined by electrophoresis on 1% (w/v) TAE agarose gel having 0.1 μ l/mL ethidium bromide (10). The amplicon were then visualized on a UV transilluminator and photographed (BioDoc-Analyse; Biometra, Goettingen, Germany).

Table 1: DNA Primer which purchased from Bioneer (Korea) company.

Primer	Sequence		Amplicon
16S rRNA	F	TCTTGTGAGAGCGGGGATA	725bp
	R	AGTTGCAGACTCCAATCCGG	
TEM-3	F	TGCATCTTTGAGCGCTCTGA	318bp
	R	CGTTTTCTGAGACGACCCCA	
TEM-72	F	TCCTTGAGAGTTTTCGCCCC	581bp
	R	CAGTGCTGCAATGATACCGC	
TEM-89	F	GGGAACCGGAGCTGAATGAA	254bp
	R	CAGTGCTGCAATGATACCGC	
TEM-160	F	CTCTAGCTTCCCGGCAACAA	149bp
	R	CAGTGCTGCAATGATACCGC	
TEM-177	F	TGATAACACTGCGGCCAACT	358bp
	R	CAGTGCTGCAATGATACCGC	
TEM-1	F	TCCTTGAGAGTTTTCGCCCC	452bp
	R	TTGTTGCCGGGAAGCTAGAG	
TEM-156	F	AGATCAGTTGGGTGCACGAG	637bp

	R	CAGTGCTGCAATGATACCGC	
--	----------	-----------------------------	--

Results:

A total of 85 clinical samples have been *Proteus* ssp collected, 69 belong to *P. mirabilis*. That isolate from different source. the highest isolation rate of *P. mirabilis* was the stool samples 19 (12.02%), ear swabs in second place in isolation 24 (9.18%), either burns samples and urine have equal proportions 16 (8.64%) from each other, while blood samples proportion 0.54% just one of isolation. The other species belonging to the genus *Proteus* isolation rate 16 (8.64%). as results showed that cultural and biochemical tests and gene 16rRNA PCR in the figure(a) belong 69 isolated to *P. mirabilis*.

Table2: Incidence of the isolated *Proteus* ssp in different clinical sample Sites.

No.(%)	Source of samples					<i>Proteus</i> ssp
	blood	burn	ear	urine	stool	
(%37.29)69	(0.54) 1	(8.64) 16	(9.18) 17	(8.64) 16	(12.02) 19	<i>P. mirabilis</i>
(%8.64)16	0(0)	(1.08)2	(2.16)4	(2.16)4	(3.24)6	<i>Other Proteus</i> ssp

$X^2 = 36.129$ *found morale differentiation in the Table2

According to the gender and age, 40 (57.97%) samples were collected from female while 29 (42.02%) samples were collected from male. the highest rate of *P. mirabilis* in the age 30-21 age increased by 34.78%, followed by the age group 31-40 years, which recorded a rate of 13.4%, as shown in the table (3). While researcher didn't agree with him (14) for the class (29-20) increased by 21.78%, and (39-30) increased by 17.82%.

Table 3 : Distribution *P. mirabilis* according to the gender and age.

Age \ Gender	Female	Male	NO%
10-1	3 (4.34)	(7.24)5	8 (11.59%)
20-11	5 (7.24)	(4.34)3	8 (11.59%)
30-21	9 (13.4)	15 (21.73)	24 (34.78%)
40-31	7 (10.14)	2 (2.89)	9 (13.04%)
50-41	5 (7.24)	2 (2.89)	7 (%10.14)
>51	11 (15.94)	2 (2.89)	13 (18.84%)
NO%	40(%57.97)	29(%42.02)	69(100%)

$X^2 = 26.41$ found morale differentiation in the Table3.



Figure(a): Ethidium bromide-stained agarose gel PCR of genes examination of *P. mirabilis* have diagnosis gene 16S rRNA. Where an DNA molecular size marker M: Marker ladder 2000-100bp Lanes No. (1--24) isolates were positive for the gene(16rRNA) gene length of 305bp. The electrophoresis performed at 60 volt for 2 hr.


In The present study results showed a higher proportion of the presence gene bla_{TEM}-177, has been in isolation for 21 bacteria *P.mirabilis* by 87.50%, followed by gene bla_{TEM}-160 as Seen in isolation for 19 *P.mirabilis* and bacteria by 59.3%, as shown in the figure(b), with the proportion of the presence of Jane bla_{TEM}-160 bacterium *P.mirabilis* record 0.85%. The resistance gene bla_{TEM}-1 has a presence in 18 *P.mirabilis* isolated by (56.25%). The proportion of gene bla_{TEM}-1 64% shown in the figure(c). While the proportion of the presence of gene bla_{TEM}-3 for *P.mirabilis* reached the lowest rate of the enzyme-resistant type bla_{TEM} and by 6.25% shown in the figure(c). But gene bla_{TEM}-89 has a presence in 10 isolates only *P.mirabilis* and increased by 31.25% shown in the figure(d). the results of the current study, the proportion of the presence of resistance gene bla_{TEM}-72 in 18 isolated and by 56.25% shown in the figure(d), as the results of the present study was that the percentage of gene bla_{TEM}-159 in 17 isolated and by 53.13% shown in the figure(e).

(b)




Figure(b): Ethidium bromide-stained agarose gel mPCR of genes examination of antibiotics Group Extended spectrum beta-lactamase genes in isolates of *P. mirabilis*. Where an DNA molecular size marker M: Marker ladder 2000-100bp Lanes No. (1-8 12-15 and 17-24) isolates were positive for the gene beta-lactamase (bla_{TEM}-177) gene length of 358bp. Lanes No. (1-8 11, 12, 15 and 17-24) isolates were positive for the gene (bla_{TEM}-160) gene length of 149bp. The electrophoresis performed at 60 volt for 2 hr.

(c)




Figure(c): Ethidium bromide-stained agarose gel mPCR of genes examination of antibiotics Group Extended spectrum beta-lactamase genes in isolates of *P. mirabilis*. Where an DNA molecular size marker M: Marker ladder 2000-100bp Lanes No. (1-10 15-17 and 19-22) isolates were positive for the gene beta-lactamase (*bla_{TEM}-1*) gene length of 452bp. Lanes No. (3 and 18) isolates were positive for the gene (*bla_{TEM} -3*) gene length of 318bp. The electrophoresis performed at 60 volt for 2 hr.

(d)



Figure(d): Ethidium bromide-stained agarose gel mPCR of genes examination of antibiotics Group Extended spectrum beta-lactamase genes in isolates of *P. mirabilis*. Where an DNA molecular size marker M: Marker ladder 2000-100bp Lanes No. (2-5 ,9,10, and 13-24) isolates were positive for the gene beta-lactamase (*bla_{TEM}-72*) gene length of 581bp. Lanes No. (2-5,9,10 and 13-17) isolates were positive for the gene (*bla_{TEM} -89*) gene length of 254bp. The electrophoresis performed at 60 volt for 2 hr.

(e)



Figure(e): Ethidium bromide-stained agarose gel PCR of genes examination of antibiotics Group Extended spectrum beta-lactamase genes in isolates of *P. mirabilis*. Where an DNA molecular size marker M: Marker ladder 2000-100bp Lanes No. (1-8 11-13 and 17-15 ,21 and 22) isolates were

positive for the gene beta-lactamase (*bla_{TEM}*-156) gene length of 637bp. The electrophoresis performed at 60 volt for 2 hr.

Table4:relationship between genes *bla_{TEM}* and source of *P.mirabilis*

Gene \ Source	Ear	Stool	Urine	Burn	Blood
<i>bla_{TEM}</i> -177	6(5.7%)	6(5.7%)	5(4.7%)	3(2.8%)	1(0.9%)
<i>bla_{TEM}</i> -160	5(4.7%)	6(5.7%)	5(4.7%)	2(1.9%)	1(0.9%)
<i>bla_{TEM}</i> -72	7(6.6%)	3(2.8%)	4(3.8%)	3(2.8%)	1(0.9%)
<i>bla_{TEM}</i> -1	7(6.6%)	5(4.7%)	4(3.8%)	2(1.9%)	1(0.9%)
<i>bla_{TEM}</i> -156	7(6.6%)	4(3.8%)	4(3.8%)	1(0.9%)	1(0.9%)
<i>bla_{TEM}</i> -89	4(3.8%)	3(2.8%)	1(0.9%)	1(0.9%)	1(0.9%)
<i>bla_{TEM}</i> -3	1(0.9%)	1(0.9%)	0(0%)	0(0%)	0(0%)
Sum	37(35.2)	28(26.6%)	23(21.9%)	12(11.4%)	6(5.7%)

$$X^2=71.321$$

*found morale differentiation in the Table4

Table5:relationship between genes *bla_{TEM}* and the age.

Gene \ Age	<i>bla_{TEM}</i> -177	<i>bla_{TEM}</i> -160	<i>bla_{TEM}</i> -1	<i>bla_{TEM}</i> -72	<i>bla_{TEM}</i> -156	<i>bla_{TEM}</i> -89	<i>bla_{TEM}</i> -3	SUM
1-10	4(3.80)	3(2.85)	2(1.90)	4(3.80)	2(1.90)	2(1.90)	--	17(70.83%)AC
20-11	5(4.76)	5(4.76)	2(1.90)	2(1.90)	4(3.80)	3(2.85)	1(0.95)	22(91.66%)AB
30-21	5(4.76)	4(3.80)	4(3.80)	5(4.76)	3(2.85)	2(1.90)	--	23(95.83%)B
40-31	5(4.76)	2(1.90)	1(0.95)	2(1.90)	1(0.95)	1(0.95)	--	12(50%)C
50-41	3(2.85)	3(2.85)	3(2.85)	2(1.90)	6(5.71)	2(1.90)	--	19(79.16%)ABC
>51	2(1.90)	2(1.90)	6(5.71)	3(2.85)	1(0.95)	1(0.95)	1(0.95)	16(66.66%)C

$$X^2=18.996$$

*found morale differentiation in the Table 5.

Table6:relationship between genes *bla_{TEM}* and the gender.

Gene \ Gender	Femle	Male	Sum
<i>bla_{TEM}</i> -177	12(11.42)	9(8.57)	21(20)
<i>bla_{TEM}</i> -160	6(5.71)	13(12.38)	19(18.09)
<i>bla_{TEM}</i> -72	8(7.61)	10(9.52)	18(17.14)
<i>bla_{TEM}</i> -1	13(12.38)	5(4.76)	18(17.14)
<i>bla_{TEM}</i> -156	11(10.47)	6(5.71)	17(16.19)
<i>bla_{TEM}</i> -89	7(6.66)	3(2.85)	10(9.52)
<i>bla_{TEM}</i> -3	1(0.95)	1(0.95)	2(1.90)
Sum	58(55.23%)	47(44.76%)	105 (100%)


$$X^2=22.271$$

*found morale differentiation in the Table 6.

Phylogenetic tree analysis to *bla_{TEM}* gene (DNA Sequencing) , Test yields 14 sample of isolates *P.mirabilis*. They took two isolation each gene of *bla_{TEM}* studied a *bla_{TEM}*-1, *bla_{TEM}*-3, *bla_{TEM}*-72, *bla_{TEM}*-89, *bla_{TEM}*-156 *bla_{TEM}*-160, *bla_{TEM}*-177 compared isolates of *P.mirabilis*

resistant of β -lactamas antibiotics with worldwide origin *P.mirabilis*.

(f)



Figure(f): Analyzed Phylogenetic tree genetic tree analysis by Mega6 program, were used genetic tree analysis of the type (Unweighted Pair Group Method) UPGMA tree Shown.

In the Figure (g) nor differences between the sequences gene bla_{TEM} local isolates *P.mirabilis* and bla_{TEM} global isolates *P.mirabilis* But there are differences between the variants of the gene TEM, example when comparing bla_{TEM}-89 record in the beginning of the Figure (g) referred to three number with a different bla_{TEM}-72, But when comparing bla_{TEM}-89 and bla_{TEM}-3 non-existent bacteria *P.mirabilis* so compared with *Klebsiella pneumoniae* because it's nearest sequence and that bacteria intestinal contain the same sequences of enzymes resistance to it from the method to acquired through plasmids passed on between the same family. So during the study recorded for the first time gene bla_{TEM}-3 in bacteria *P.mirabilis* at the local level in the city of Diwaniyah, in the global level in the bacteria. And do not record this gene in the bacteria in GenBank. bla_{TEM}-3 presence *P.mirabilis* this gene in two isolation only, one the source ear infection while the second isolation from a patient have 18 year old person with a case of bacteremia due to bacteria *P.mirabilis* in the city of Diwaniyah. different this gene for bla_{TEM}-89 and by 2.88 the first isolation and by 2.91 second in isolation either standard isolation by 7.69. This shows that the absence of a new and dangerous types of the community in the city of Diwaniyah and bacteria *P.mirabilis* despite being seen everywhere in hospitals in Diwaniyah and although resistant multiple to antibiotics and but do not exposed to mutations in the genes of resistance to antibiotics treatment and reducing infection.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
1. Proteus mirabilis beta-lactamase TEM-89 gene isolate No.1																					
2. Proteus mirabilis beta-lactamase TEM-89 gene isolate No.2	0.00																				
3. Proteus mirabilis TEM-89 beta-lactamase (TEM-89) gene (AY039040.1)	0.00	0.00																			
4. Proteus mirabilis beta-lactamase TEM-72 gene isolate No.1	0.01	0.01	0.01																		
5. Proteus mirabilis beta-lactamase TEM-72 gene isolate No.2	0.01	0.01	0.01	0.00																	
6. Proteus mirabilis beta-lactamase TEM-72 (blaTEM-72) gene (AF157413.1)	0.01	0.01	0.01	0.00	0.00																
7. Proteus mirabilis beta-lactamase TEM-3 gene isolate No.1	2.42	2.42	2.88	2.82	2.82	2.82															
8. Proteus mirabilis beta-lactamase TEM-3 gene isolate No.2	2.47	2.47	2.91	2.85	2.85	2.85	0.00														
9. Klebsiella pneumoniae beta-lactamase TEM-3 (blaTEM-3) gene (X64523.1)	3.56	3.53	7.69	4.78	4.77	7.70	0.00	0.00													
10. Proteus mirabilis beta-lactamase TEM-177 gene isolate No.1	0.02	0.02	0.02	0.01	0.01	0.01	2.69	2.72	3.45												
11. Proteus mirabilis beta-lactamase TEM-177 gene isolate No.2	0.02	0.02	0.02	0.01	0.01	0.01	2.70	2.73	3.49	0.00											
12. Proteus mirabilis beta-lactamase TEM-177 (blaTEM-177) gene (FN652295.1)	0.02	0.02	0.01	0.01	0.01	0.01	2.79	2.83	7.04	0.00	0.00										
13. Proteus mirabilis beta-lactamase TEM-160 gene isolate No.1	0.01	0.01	0.01	0.01	0.01	0.01	2.79	2.94	4.93	0.02	0.02	0.02									
14. Proteus mirabilis beta-lactamase TEM-160 gene isolate No.2	0.01	0.01	0.01	0.01	0.01	0.01	2.80	2.95	4.77	0.02	0.02	0.02	0.00								
15. Proteus mirabilis beta-lactamase TEM-160 (blaTEM-160) gene (EF136377.1)	0.00	0.00	0.00	0.01	0.01	0.00	2.79	2.83	6.92	0.01	0.01	0.01	0.00	0.00							
16. Proteus mirabilis beta-lactamase TEM-156 gene isolate No.1	0.01	0.01	0.01	0.01	0.01	0.01	2.85	2.89	5.43	0.02	0.02	0.02	0.01	0.01	0.01						
17. Proteus mirabilis beta-lactamase TEM-156 gene isolate No.2	0.01	0.01	0.01	0.01	0.01	0.01	2.86	2.89	5.38	0.02	0.02	0.02	0.01	0.01	0.01	0.00					
18. Proteus mirabilis beta-lactamase TEM-156 (blaTEM-156) gene (AM941159.1)	0.01	0.01	0.01	0.01	0.01	0.01	2.86	2.89	8.29	0.02	0.02	0.01	0.01	0.01	0.01	0.00	0.00				
19. Proteus mirabilis beta-lactamase TEM-1 gene isolate No.1	0.00	0.00	0.01	0.01	0.01	0.01	2.93	2.93	4.65	0.02	0.02	0.01	0.00	0.00	0.01	0.01	0.01	0.01			
20. Proteus mirabilis beta-lactamase TEM-1 gene isolate No.2	0.00	0.00	0.01	0.01	0.01	0.01	2.93	2.93	4.65	0.02	0.02	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.00		
21. Proteus mirabilis beta-lactamase TEM-1 gene (AY729027.1)	0.01	0.01	0.01	0.01	0.01	0.01	2.88	2.91	8.28	0.02	0.02	0.01	0.01	0.01	0.01	0.00	0.00	0.01	0.00	0.00	0.00

Figure 11-4: measure the genetic differences in the genetic sequence of bases between the genes of beta-lactamase TEM genes in the local *P.mirabilis* in Diwaniyah city and compared with the beta-lactamase TEM genes in the *P. mirabilis* recorded in the world.

Discussion

According to Isolation and Identification of *P. mirabilis* in this study agree (11) with a percentage 45%. Also The result of current study didn't agree with (12). According to the gender that nearest with (13) and didn't agree with (14) belong the infection - to bacterial may be due to the fact that extreme ages (children and seniors) and especially the elderly who that have from chronic diseases and those taking medications immunosuppressive, they are more exposed to infection because of a weakened immune system for these age groups compared to other age groups (15) .

The enzymes bla_{TEM} of more enzymes beta-lactamase prevalent that goes back to Class A, which consists of three main groups of enzymes (, bla OXA, blaCTX-M, blaSHV blaTEM), but is blaTEM of the main kinds and these enzymes most likely found in *E.coli* and *K.pneumoniae* and *P.mirabilis* (17). gene bla_{TEM} -160 in this study are consistent with the researcher (18), with the proportion of the presence of blaTEM -160 bacterium *P.mirabilis* record 0.85%. The resistance gene bla_{TEM} -1 has converged these results with the results of with (19) .In another study in the South China conducted by (20) recorded bla_{TEM} -1 at a higher rate than the current study 78%. gene blaTEM-3 for *P.mirabilis* record the gene bla_{TEM} -3 in 1987 as the first different type of enzymes the blaTEM where he owns an increase in potency against cephalosporin antibiotics broad-spectrum (21). But gene bla_{TEM}-89 consistent with the researcher (22). gene blaTEM-72 as the person researcher (23) The results of the DNA Sequencing analysis, there is a clear Identify of isolates of *P.mirabilis* blaTEM local isolates in Diwaniyah city with worldwide origin *P.mirabilis* compared with the rest of the other species that appear in the .

REFERENCES:

1. Armbruster, C. E. and Mobley, H. L. 2012. Merging mythology and morphology: the multifaceted lifestyle of *Proteus mirabilis*. *Nature Reviews Microbiology*. 10(11), pp: 743-754

2. **Jacobsen, S. M. & Shirtliff, M. E. (2011).** *Proteus mirabilis* biofilms and catheter-associated urinary tract infections. *Virulence* 2, 460–465. *Iran. EMHJ*, 18(2).
3. **Pellegrino, R.; Scavone, P.; Umpiérrez, A.; Maskell, D.J. and Zunino, P. (2013).** *Proteus mirabilis* uroepithelial cell adhesin (UCA) fimbria plays a role in the colonization of the urinary tract. *J. Pathol. Dis.*, 2(67):104-107.
4. **Mollenkopf, D. (2012).** Epidemiology of CTX-M cephalosporinase – bearing *Escherichia coli* and *Salmonella* spp isolated from US livestock .M.Sc .Thesis, college of science ,Ohio state University .
5. **Dhillon R, Clark J (2011).** ESBL: A clear and present danger? *Crit. Care Res. Pract.* 2012 (2012):1-11.
6. **Singh, V.; Sharma, L.; Kanta, R.; Sharma, S.; Chauhan, S. and Chauhan, P. K. (2010).** Antimicrobial activity of *Proteus* bacteria isolated from raw milk of Paonta Valley. *Int. J. Phar. Bio Sci.* 1(2):1-4.
7. **D'Andrea MM, Literacka E, Zioga A, et al. (2011).** Evolution and spread of a multidrug-resistant *Proteus mirabilis* clone with chromosomal AmpC-Type Cephalosporinases in Europe. *Antimicrob Agents Chemother.* 55(6):2735-42.
8. **Meervenne EV, Coillie EV, Kerckhof FM, et al. (2012)** Strain specific transfer of antibiotic resistance from an environmental plasmid to foodborne pathogens. *Journal of Biomedicine and Biotechnology*, ID 834598, doi:10.1155/2012/834598.
9. **Sambrook, J.; Fritsch, E.F. and Maniatis, T. (1989).** Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
10. **Bartlett, J.M.S. and Stirling, D. (1998).** PCR Protocols: Methods in molecular biology. 2nd. Humana Press Inc. Totowa.
11. **Zuhir , R; Alaubydi, M. A. S. (2016).** Extraction and Partial Purification of Lipopolysaccharide from Clinical *Proteus mirabilis* Isolate and Compared with Standard Bacteria. *Iraqi Journal of Science*, 2016, Vol. 57, No.1C, pp: 599-608.
12. **Kwiecinska-Pirog , J ; Skowron ,K ; Bartczak ; and Gospodarek-Komkowska, E. (2016).** The Ciprofloxacin Impact on Biofilm Formation by *Proteus Mirabilis* and *P. Vulgaris* Strains . *Jundishapur J Microbiol.* 9(4):e32656.
13. **Senthamarai ,S. Sivasankari, S.; Anitha ,C.; Kumudavathi ,MS.; Amshavathani ,SK.; Venugopal ,V.; and Thenmozhi Valli, P R. (2015).** A study on the antibiotic susceptibility pattern of *Proteus* spp among various samples. *IJAPBC* – Vol. 4(2): 2277 – 4688.
14. **Pal, N.; Sharma, N.; Sharma, R.; Hooja, S.; and K Maheshwari , R. (2014) .** Prevalence of Multidrug (MDR) and Extensively Drug Resistant (XDR) *Proteus* species in a tertiary care hospital, India. *Int. J. Curr. Microbiol. App. Sci* 3(10) 243-252.
15. **Brooks , G.F. ; Carroll , K. C. ; Butel , J.S. and Morse , S. A. (2007) .** Jawetz, Melnick, Adelberg's Medical microbiology. 24th ed. *McGraw-Hill com.* PP. 174.

- 16. Guilfoile, G.(2007).** Deadly disease and epidemics Antibiotic-Resistant Bacteria. Chelser House publisher , *World Health organization*. P:10-119.
- 17. Sharma, J.; Sharma, M. and Roy, P.(2010).** Detection of TEM and SHV genes in *Escheria coli* and *Klebsiella pneumoniae* isolates in a tertiary care hospital from India, *Indian J. Med. Res.*, 132(3):332-336.
- 18. Arago'n, L. M.; Mirelis ,B.; Miro', E.; Mata,C. ; Go'mez, L.; Rivera,A. ; Coll1,P.; and Navarro1,F.(2008).** Increase in b-lactam-resistant *Proteus mirabilis* strains due to CTX-M- and CMY-type as well as new VEB- and inhibitor-resistant TEM-type b-lactamases. *Journal of Antimicrobial Chemotherapy* ,61: 1029–1032.
- 19. Jones, C. H.; Tuckman, M.; Keeney,D.; Ruzin,A.; and Bradford, P.A. (2009) .** Characterization and Sequence Analysis of Extended-Spectrum- β -Lactamase-Encoding Genes from *Escherichia coli*, *Klebsiella pneumoniae*, and *Proteus mirabilis* Isolates Collected during Tigecycline Phase 3 Clinical Trials . *ANTIMICROBIAL AGENTS AND CHEMOTHERAPY* .Vol 53(2) p: 465–475.
- 20. Pellegrino. R.; Scavone, P.; Umpiérrez, A.; Maskell, D.J. and Zunino, P.(2013).** *Proteus mirabilis* uroepithelial cell adhesin (UCA) fimbria plays a role in the colonization of the urinary tract. *J. Phatho. Dis.*, 2(67):104-107.
- 21. Bradford, P.A. (2001).** Extended-spectrum β -Lactamase in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin. Microbiol. Rev.* 14:933-951.
- 22. NEUWIRTH,C.; MADEC,S.; SIEBOR,E .; PECHINOT,A.; DUEZ, J.; PRUNEAUX,M.; FOUCHEREAU-PERON,M.; KAZMIERCZAK,A. AND LABIA, R.(2001).** TEM-89 β -Lactamase Produced by a *Proteus mirabilis* Clinical Isolate: New Complex Mutant (CMT 3) with Mutations in both TEM-59 (IRT-17) and TEM-3. *ANTIMICROBIAL AGENTS AND CHEMOTHERAPY*, 45(12) :3591–3594.
- 23. PERILLI, M.; SEGATORE,B.; MASSIS,M. R. D.; RICCIO, M. L.; BIANCHI,C.; ZOLLO,A.; ROSSOLINI,G.M.; AND AMICOSANTE, G (2000).** TEM-72, a New Extended-Spectrum b-Lactamase Detected in *Proteus mirabilis* and *Morganella morganii* in Italy. *ANTIMICROBIAL AGENTS AND CHEMOTHERAPY*, 44(9) :2537–2539 .