



**Republic of Iraq**  
**Ministry of Higher Education and**  
**Scientific Research**  
**University of Al-Qadisiya**  
**College of Veterinary Medicine**



ISSN: 1818 – 5746

# **Al-Qadisiya Journal of Veterinary Medicine Sciences**

SCIENTIFIC JOURNAL PUBLISHED BY THE  
COLLEGE OF VETERINARY MEDICINE  
UNIVERSITY OF AL-QADISIYA

**Iraq, Diwania, College of Veterinary Medicine**  
email: [qjvms@yahoo.com](mailto:qjvms@yahoo.com) , [vmjou@qu.edu.iq](mailto:vmjou@qu.edu.iq)

Chief Editor. Tel. 0780 1786143  
Editorial Secretary. Tel. 0780 3105771

**Vol. 14 No. 1 2015**

**Republic of Iraq**  
**Ministry of Higher Education and Scientific Research**  
**University of Al-Qadisiya**  
**College of Veterinary Medicine**

## **Al-Qadisiya Journal of Veterinary Medicine Sciences**

**Is half annual scientific peer reviewed journal published by the  
College of Veterinary Medicine University of Al-Qadisiya**

### **Editorial Board**

**Editor-in- Chief: Prof. Dr. Thaier Alwan Abid**

**Editorial Secretary: Asst. Prof. Dr. Qassim Haleem Kshash**

### **Editors**

Prof. Dr. Amina N. Thuwani

Prof. Dr. Alia Y. Yaqob

Prof. Dr. Salah M. Hassan

Prof. Dr. Baraa N. Abdullah

Prof. Dr. Mohammad J. Eesa

Prof. Dr. Rahman K. Mohsen

Asst. Prof. Dr. Najah H. Hassan

Asst. Prof. Mr. Kareem N. Tahir

**Al-Qadisiya Journal of Vet. Med. Sci. is printed and organized by**

Asst. Prof. Dr. Eman. F. Al- Baghdady

Lecturer Dr. Baha. A. Abdul-Hussien

Lecturer Dr. Mansor. J. Ali

Technical Manager Basema. E.M. Al-Sa'dy

### **Staff of the editorial office**

Asst. Prof. Dr. Jenan A. S. Alhussaini

Lecturer Dr. Dhafer R. H. Alfetly

### **Directed by**

Dr. Thaier Alwan Abid

Mrs. Basema E.M. Al-Sa'dy

Articles published in this journal are available as pdf in [qu.edu.iq/vmjou](http://qu.edu.iq/vmjou), and Iraqi Academic Scientific Journals (IASJ).

**Vol. 14 No. 1 2015**

Contents

Research articles	page
<b>Detection of Giardiasis in apparently healthy cattle by using direct ELISA technique</b> <i>Heba Shehab Ahmed, and Noaman Naji A'aiz</i> .....	1
<b>Study the effect of green tea extract on lipid profile in nitrate treated rabbits</b> <i>Hayder G. A. AL-fatlawi and Hayder A. N. AL- Zamely</i> .....	4
<b>Estimation the alkaline phosphates level in partial hepatectomy rabbits</b> <i>Humam H. Nazht, Nabeel Abdul-Rahman AL-Keledar and Abeer Ahmed Majeed</i> .....	10
<b>Prevalence of coccidiosis in local breed rabbits (<i>Oryctolagus cuniculus</i>) in Baghdad province</b> <i>Athraa Toma Khider, Haider M. A. Al-Rubaie and Farouk Jummah Khalil</i> .....	15
<b>Rapid detection of infectious bronchitis virus in broilers in Al-Diwaniya governorate by using Real-Time reverse transcriptase Polymerase Chain Reaction</b> <i>Abbas Hadi Jasim Al-Mahmoudi and Nafea S. Jasim</i> .....	22
<b>Some molecular and sero – prevalence study of <i>Anaplasma marginale</i> in cattle in Wassit province</b> <i>Ola Abd-El Hussain Agaar and Ghyda'a Abbas Jassem</i> .....	27
<b>Rapid detection of <i>Yersinia enterocolitica</i> by using Real-Time PCR technique in some types of foods in Al-Qadisiya province</b> <i>Kadhim H. Abbas</i> .....	34
<b>The effect of running at many speeds on some synovial fluid constituents in Iraqi Arabian horses</b> <i>Amir Ibrahim Towfik</i> .....	39
<b>Study the pathogenicity of <i>Enterobacter cloacae</i> in rats that isolated from diarrhetic buffalos calves in Babylon province</b> <i>Abdul-kareem Salman Al-Yassari</i> .....	44
<b>Behavioral, hormonal, and testicular histological study to evaluate the effect of season on mating in local breed free housing tom cats in Al-Diwaniya city</b> <i>Ali H. Jaber AL-bdeery</i> .....	49
<b>RAPD-PCR based genetic variation of <i>Candida albicans</i> of animal and human origin</b> <i>Hayder Mahmood Ali Samaka</i> .....	54
<b>Prevalence of toxoplasmosis in renal infections patients in Al-Muthanna province/ Iraq</b> <i>Mohenned A. Hemza AlSaadawi, and Mansoor Jadaan Ali Alkhaled</i> .....	58
<b>A comparison of gravid uterine parameters of local breed ewes between single and twin pregnancies in different gestational stages</b> <i>Hella J. Al-Fatlawy</i> .....	61
<b>The effect of pregnancy and lactation on blood biochemical and immunological values in ewes after enterotoxaemia vaccination</b> <i>Basim H. Abed Ali</i> .....	65
<b>Hematological and biochemical changes following Ivermectin treatment in mange infested goats</b> <i>Batool Kathem Habeeb</i> .....	69

<b>Prevalence of <i>Rota virus</i>, <i>Escherichia coli</i> O157:H7 and <i>Cryptosporidium spp.</i> in feces of buffalo in Babil governorate</b> <i>Amanee Mohammad Radhy</i> .....	75
<b>Comparative anatomical and histological study of the esophagus of local adult male and female homing pigeon (<i>Columba livia domestica</i>)</b> <i>Khalid Hadi Kadhim, and Ahmed Adeeb Mohamed</i> .....	80
<b>Effects of sub lethal doses of endosulfan on histopathological and biochemical parameters of common carp (<i>Cyprinus carpio</i>)</b> <i>Mohanad obayes abdullah AL-jubouri</i> .....	88
<b>Laparoscopic colotomy suture using clips and Connell techniques in goats: A comparative study</b> <i>Rafid H. Farman, and Mohammad J. Eesa</i> .....	95
<b>Study the association between <i>Enterobius vermicularis</i> infection and enuresis among children in Al-Najaf city</b> <i>Huda A S Al-Shadoo</i> .....	104
<b>Effect of <i>Punica granatum</i> rinds ethanolic extract on healing of fungated wounds in rabbits</b> <i>Muna H. AL-Saeed, Rasha M. Othman and Arwa H. AL-Saeed</i> .....	109
<b>Histomorphological and morphometrical comparative study of the kidney between Quail (<i>Conturnix coturni</i>) and Green-winged Teal (<i>Anas crecca</i>) according to their environment type</b> <i>Ali faidh Baragoth</i> .....	118
<b>Detection of hemolysin virulence factor gene of Enterohemorrhagic <i>Escherichia coli</i> (EHEC) isolated from feces of infected sheep by using Polymerase Chain Reaction Technique</b> <i>Saba Aboud Ali</i> .....	127
<b>Evaluation the clinical effects of neuroleptanalgesia (Remifentanil-Acepromazine, Remifentanil-Xylazine, and Remifentanil-Midazolam) during intubation and some minor surgical operations in dogs</b> <i>Samir Aoda Jaffar and Ayad Abed-Al-Jabbar Amin</i> .....	132



## Rapid detection of *Yersinia enterocolitica* by using Real-Time PCR technique in some types of foods in Al-Qadisiya province

Kadhim H. Abbas

Coll. of Vet. Med. / Univ. of Al-Qadisiya

email: [kadhimhealth@gmail.com](mailto:kadhimhealth@gmail.com)

(Received 13 April 2014, Accepted 22 May 2014)

### Abstract

The present study was carried out for direct and rapid detection of pathogenic *Y. enterocolitica* in some types of foods such as (raw milk, local sweet cheese and minced beef meat) exist in local different markets of Al-Qadisiya province. A total of 192 samples were collected randomly from markets of Al-Qadisiya province involved 79 raw milk samples, 58 local sweet cheese samples and 55 minced beef meat samples. Real-Time PCR technique based SYBER Green dye were used for detection of this bacteria by amplifying invasion locus protein (*ail*) gene which is found especially in *Y. enterocolitica*, The results revealed that positive samples for this (*ail*) gene were 49 samples out of 79 examined raw milk (62%), also 6 samples out of 58 samples observed in local sweet cheese (10.3%) and 49 samples out of 55 samples of minced beef meat (89%). A total of positive samples were 104 out of 192 examined samples, samples of minced beef meat recorded highly significant contamination ( $p < 0.05$ ) as compared samples of raw milk which occupied second rank in contamination with pathogen and then samples of local sweet cheese which recorded less degree of contamination. The Real-time PCR technique was very specific and efficient for detection *Y. enterocolitica* in variety foods, when compared with culture method which have many problems.

**Keywords:** *Y. enterocolitica*, Real-Time PCR, *ail* gene, raw milk, sweet cheese, minced beef meat.

### الكشف السريع عن اليرسينيا القولونية المعوية باستخدام تقنية تفاعل البوليمر السلسلي المؤلف في بعض انواع الغذاء في محافظة القادسية

كاظم حسن عباس

كلية الطب البيطري/ جامعة القادسية

### الخلاصة

اجريت الدراسة الحالية للكشف المباشر والسريع عن اليرسينيا القولونية المعوية في بعض انواع الاغذية الموجودة في الاسواق المحلية في محافظة القادسية كالحليب الخام، الجبن الحلو المحلي، ولحم العجل المفروم جمعت 192 عينة عشوائياً من اسواق محافظة القادسية شملت 79 عينة حليب خام و58 عينة جبن حلو محلي و55 عينة لحم عجل مفروم. استخدمت تقنية تفاعل البوليمر السلسلي المؤلف والمعتمد على صبغة السايبر غرين لكشف هذه البكتيريا وذلك بمضاعفة بروتين الغزو الموضوعي للـ (*ail*) جين والذي يتواجد حصرياً في بكتيريا اليرسينيا القولونية المعوية. بينت النتائج ان العينات الموجبة للـ (*ail*) جين كانت 49 عينة من 79 عينة من الحليب الخام المفحوصة (62%) و 6 عينات من 58 عينة من عينات الجبن الحلو المحلي المفحوصة (10.3%) وفي 49 عينة من 55 عينة من عينات لحم العجل المفروم المفحوصة (89%) ، مجموع العينات الموجبة كان 104 من 192 عينة مفحوصة. سجلت عينات لحم العجل المفروم ارتفاعاً معنوياً ( $p < 0.05$ ) في التلوث مقارنة بعينات الحليب الخام والتي جاءت بالمرتبة الثانية تلوئاً بالبكتيريا ومن ثم عينات الجبن الحلو المحلي والتي كانت اقل تلوئاً. وأظهرت النتائج ان تقانة تفاعل البوليمر السلسلي المؤلف تعد طريقة دقيقة جداً وفعالة في تشخيص اليرسينيا القولونية المعوية في الاغذية المتنوعة مقارنة مع طريقة الزرع الجرثومي الاعتيادية والتي لها معوقاتهما العديدة. الكلمات المفتاحية: اليرسينيا القولونية المعوية ، تقنية تفاعل البوليمر السلسلي ، جين الـ *ail* ، الحليب الخام ، الجبن الحلو ، لحم العجل المفروم.

## Introduction

*Y. enterocolitica* is a species of gram-negative coccobacillus-shaped bacterium classified belonging to Enterobacteriaceae, a pathogenic strain is characterized by a specific in the epidemiological mechanism of infections related with consumption of contaminated foods and associated with food borne disease called Yersinosis which characterized by diarrhea, mesenteric lymphadenitis, terminal ileitis, arthritis and septicemia (1, 2, 3, 4). The foods from animal origin play an important role in the transmission of *Yersinia spp.* (5, 6, 7). Pathogenic strain of *Yersinia* classified as psychrophilic bacteria because its ability to growing and multiplication in low-temperature of storage which used for keep of perishable food products. These reveal that the used of chilled and frozen storage of foods are potential reservoirs and sources of infection and may cause transmission of pathogenic *Yersinia spp.* (8, 9). The pathogenic potential of *Y. enterocolitica* is determined by complex interaction of chromosomal and plasmid genes (10). The presence of a 70 kb plasmid (pYV, plasmid *Yersinia* virulence) and specific chromosomal determinants (genes *ail*, *yst*, etc.) are necessary for the full expression of pathogenic potential (10, 11). Real-Time PCR is the developed method up to now based on *in vitro* amplification and has a potential for rapid and selective detection of microorganisms during the last years there is a significant progress in development of Real-Time PCR aiming at the quantitation of bacterial load in different complex samples, its principle based on detection of a fluorescent signal which is proportional to the number of amplicons in the tested sample, the main goal of the recent study is the optimization of a Real-Time PCR protocol based on the *ail* gene in order to quantify *Y. enterocolitica* (3, 12). This will contribute to the fast and quantitative detection of *Y. enterocolitica* in complex samples like meat, milk products and feces (1, 13). Detection of *Y. enterocolitica* in daily consummated food are important for consumer's health because inadequate healthy processing during preparation of

these food materials (14). The aim of the present study was rapid detection of *Y. enterocolitica* by using Real-time PCR technique in some types of foods in Al-Qadisiya province.

## Materials and methods

### Samples collection

A total of 192 Samples of raw milk, local sweet cheese, and minced beef meat were collected randomly from different local markets in Al-Qadisiya province. The samples were collected in 25ml sterile containers and transported to laboratory and stored in a refrigerator until use for genomic DNA extraction accept raw milk which examined three hours after collection .

### Genomic DNA extraction

Bacterial genomic DNA was extracted from milk, cheese, minced beef meat according method described by (15) by using (Genomic DNA extraction kit. Bioneer. Korea) with some modification. 1ml aliquot of milk was centrifuged at  $6,000 \times g$  for 10 min, and then the clear portion was pipette and discarded. The remaining milk solids and butter fat were used for DNA extraction, while, the cheese 250mg directly used for extraction. Whereas, 200mg from minced meat was homogenized by tissue lysis buffer that used for DNA extraction. The others steps were done according to kit instruction using DNA purification spin column. After that, the purified DNA eluted in elution buffer provided with kit and the extracted DNA was checked by Nano-drop spectrophotometer, then the DNA store at  $-20^{\circ}\text{C}$  until used for prepared of qPCR master mix.

### Real-Time PCR

Real-Time PCR technique was used to detection of *Y. enterocolitica* food borne bacteria using SYBER green dye based amplification of attachment invasion locus protein (*ail*) gene in *Y. enterocolitica* from milk, cheese, and minced meat. This technique was carried out according to method described by (3, 12). Real-Time PCR primers were designed by this study from conserved region attachment invasion locus



protein (*ail*) gene (GenBank: JX972144.1) using NCBI-GenBank data base and primer3 plus online, and these primers were provided by Bioneer Company. Korea as showed in table (1).

**Table (1): structure of primer**

Primer		Sequence	Amplicon
<i>ail</i>	F	GGATGCAATTCAAC CCTACTTC	98bp
	R	TGCACCAAGTATCCA AGTGC	

The Real-Time PCR amplification reaction was done by using (AccuPower™ 2X Green star qPCR master mix kit, Bioneer. Korea) and the qPCR master mix were prepared for each sample according to company instruction as in table (2).

**Table(2) Composition of qPCR master mix**

qPCR master mix	Volume
DNA template	3µL
2X Green star master mix	25µL
<i>ail</i> Forward primer (10pmol)	1.5µL
<i>ail</i> Reverse primer (10pmol)	1.5µL
DEPC water	19µL
Total volume	50µL

These qPCR master mix reaction components that mentioned in table were placed in sterile white qPCR strip tubes and transferred into Exispin vortex centrifuge for 3minutes, the place in MiniOpticon Real-Time PCR system and applied the following thermo cycler conditions as in the table (3).

**Table (3): Thermo cycler conditions**

qPCR step	Temperature	Time	Repeat cycle
Initial Denaturation	95 °C	3 minutes	1
Denaturation	95 °C	10 sec	40
Annealing\ Extension	58 °C	30 sec	
Detection(scan)			
Melting	60-95°C	0.5 sec	1

### Statistical analysis

The data were analyzed using Chi-square ( $p < 0.05$ ) to detect significant differences among each groups (16).

### Results

The results of Real-Time PCR assay for detection of contamination with *Y.*

*enterocolitica* shown from 192 examined samples 104 were positive with percentage (54.1%), this results distributed to 49 positive raw milk samples from 79 examined samples with percentage (62%), while 6 samples were positive from 58 examined samples of sweet local cheese with percentage (10.3%), and 49 samples were positive from 55 examined minced beef meat with percentage (89%) (Table 4).


**Table (4): The occurrence of pathogenic *Y. enterocolitica* in some types of foods.**

Items	No. of tested samples	No. of positive samples	%
raw milk	79	49	62 <sup>b</sup>
Sweet local cheese	58	6	10.3 <sup>c</sup>
Minced beef meat	55	49	89 <sup>a</sup>
Total	192	104	54.1

<sup>abc</sup> The different superscript letters in the column refer to the significant differences ( $p < 0.05$ ),  $\chi^2$  cal. = 59.633.


The contaminated samples of minced beef meat recorded highly significant differences ( $p < 0.05$ ) as compared to other examined samples.

The direct detection of *Y. enterocolitica* by Real-Time PCR based SYBR Green dye was appeared highly sensitive and specific for amplification of attachment invasion locus protein (*ail*) gene and the specificity was determined by dissociation curve (Melt Curve). Where the positive amplification product samples were showed specific amplification at melt peak mainly at (Tm: 80°C) without primer dimer or nonspecific products (Fig.1,2).



**Fig. (1): Amplification plot of *Y. enterocolitica* attachment invasion locus protein (*ail*) gene by Real-Time PCR in some type of foods.**

**Fig. (2): Melt curve of *Y. enterocolitica* attachment invasion locus protein (*ail*) gene by Real-Time PCR in some type of foods. Shows the specificity of gene amplification at (T<sub>m</sub>: 82.5°C).**



## Discussion

Real-time PCR method is a specific and rapid technique for rapid detection of *Y. enterocolitica* in different foods (1, 17). The designed primers of attachment invasion locus protein *ail* gene in this study were chosen as an excellent target for pathogenic *Y. enterocolitica* because their limited virulence strains of *Y. enterocolitica* rather than nonpathogenic *Yersinia spp.* these results are in accordance with (10) who attributed that to heterogeneity within the bacterial population for the presence of the virulence plasmid and associated with human pathogenicity. (11,18) refer to the presence of pathogen which related with *ail* gene was recommended in Europe and America and other world countries which associated with a pathogenicity of human. Conventional PCR does not differentiate between *Y. enterocolitica* and *Y. Pseudotuberculosis* but only Real time PCR method is developed for detection to *ail* gene which is found in pathogenic *Y. enterocolitica* as pointed by (2,13). Also (6) refers to the contamination of variety of foods occurs at high levels and this need a highly specific and rapid method for detection of indicator pathogen. In addition to that, (18) reaches to, there are many problems during culturing of this *Y. enterocolitica* because doesn't not well

compete with other pathogenic species. High numbers of other species of Enterobacteriaceae inhibit growth of *Y. enterocolitica* has found by (10). Highly presence of bacteria in raw milk resulted from bad handling during milking and storage of milk in contaminated utensils where the bacteria transfer from the farm to the market and without use heat treating, similar results were obtained by (1). In local sweet cheese recording low level of contamination, this result can be attributed to heat treating of raw milk during processing of making cheese, contamination happens during handling but that will not decrease the risk of this food because of eating directly; this accordance with (3). Highly contamination of minced beef meat samples resulted from high contamination in slaughterhouse and contaminated water which was used for washing of carcasses in addition that bad handling with meat in the local markets. This results similar to that reported by (7,12). In public health field when the indicator bacteria as *Y. enterocolitica* was detected in any type of foods this means that food unfit for human consumption.

## References

- 1-Asker A A, Abd El Aal S F A, Amer I H (2013) Prevalence of virulent *Yersinia enterocolitica* in subclinical mastitic cow milk in Sharkia Governorate, Egypt. Life Science Journal, 10(3): 1285-1294.
- 2-Fukushima H, Shimizu S, Inatsu Y (2011) *Yersinia enterocolitica* and *Yersinia pseudotuberculosis* detection in food. j. of Pathology., 10(40): 1-9.
- 3-Lambertz S T, Nilsson C, Hallanvuori S, Lindblad M (2008) Real-Time PCR method for detection of pathogenic *Yersinia enterocolitica* in food. Appl. Environ. Microbiol., 74(19): 6060-6067.
- 4-Lamps I W, Havens J M, Gilberch L J, Dube P H, Scott M A (2006) Molecular biogrouping of Pathogenic *Yersinia enterocolitica*. Am. J. Clin. Pathol., 125: 658-664.
- 5-Falcao J P, Falcao D P, Pitondo-Silva A, Malaspina A C, Brocchi M (2006) Molecular typing and virulence markers of *Yersinia enterocolitica* strains from human, animal and food origins isolated between 1968 and 2000 in Brazil. Med. Microbiol., 11: 1539-1548.
- 6-Homtaiz H, Rahimianian M D, Dehkordi F S (2013) Identification and characterization of *Yersinia enterocolitica* isolated from raw chicken meat based on molecular and biological technique. J. Appl. Res., (22)1: 137-145.



- 7-Sulakvelidze A, Dalakishvili K, Barry E, Wauters G, Robins-Brown R, Imnadze P, Morris J G (1996) Analysis of clinical and environment *Yersinia* isolates in republic of Georgia. *J. Clin. Microbiol.*, 4(9): 2325-2327.
- 8-Ilieve M V, Najenski H M, Werbruck H, Herman I, Coillie E V (2008) Optimization of Real-time PCR Protocol for Detection of pathogenic *Yersinia enterocolitica* strains. *Bulgarian J. of Vet. Med.*, 11(3): 179-184.
- 9-Kapperud G (1991) *Yersinia enterocolitica* in food hygiene. *Int. J. of Food Microbiol.*, 12(1): 53-65.
- 10-Thoerner P, Bin Kingombe C I, Bissig-Choisat B, Wassenaar T M, Frey J, Jemmi T (2003) PCR Detection of Virulence genes in *Yersinia enterocolitica* and *Yersinia pseudotuberculosis* and Investigation of Virulence Gene Distribution. *Appl. Environ. Microbiol. March.*, 69(3): 1810-1816.
- 11-Simonova J, Vazlerova M, Steinhauserova I (2007) Detection of Pathogenic *Yersinia enterocolitica* serotype O:3 by biochemical, serological, and PCR methods. *Czech Food Science.*, 25(4): 214-220.
- 12-Tudor L, Togoe I, Ilie L I, Tudor L, Mitranescu E (2010) The detection of *Yersinia enterocolitica* strains from waste Waters using Real-Time PCR technique. *Lucrari Stintifice medicina Veterinara.*, XLIII (1):237-242.
- 13-Zheng H, Wang J, Sun Y, Jiang B (2007) Clinical isolation and characterization of *Yersinia enterocolitica* in china using Real-Time PCR and culture method. *Digestion.*, 75: 199-204.
- 14-Fredriksson-Ahomaa M, Korkeala H (2003) Low occurrence of pathogenic *Yersinia enterocolitica* in clinical, food, and environmental samples: a methodological problem. *Clin. Microbiol. Rev.*, 16(2): 220-229.
- 15-Streevatsang S, bookout J B, Ringpis F, Perumaala V S, Ficht T A, Adama L G, Hagius S D, Elizer P H, Kumar G K, Rajasekher M, Isloor S, Barathur R (2000) Multiplex Approach to Molecular Detection of *Brucella abortus* and/or *Mycobacterium bovis* Infection in Cattle . *J. Clin. Microbiol.*, 38: 2602-2610.
- 16-Petrei A, Watson P (2004) Statistics for veterinary and animal science. illustration prepared by alexander hunte. printed and bounded in Great Britain. By T.J. Inter. Ltd, padstow Cornwall.
- 17-Lambertz S T, Danielsson-Tham M L (2005) Identification and characterization of pathogenic *Yersinia enterocolitica* Isolates by PCR and Pulsed-field Gel Electrophoresis. *Appl. Environ. Microbiol.*, 71(7): 3674-3681.
- 18-Bhaduri S, Wesley I V, Bush E J (2005) Prevalence of Pathogenic *Yersinia enterocolitica* Strains in Pigs in the United States. *Appl. Environ. Microbiol.*, 71(11): 7117-7121.