

Ministry of Higher Education And Scientific Research
University of Al-Qadisiyah
College of Computer Science & Information Technology
Department of Mathematics



Hyers-Ulam Stability Of a polynomial Equation

A research Submitted to the Department of Mathematics/College
of Computer Science & Information Technology / Al-Qadisiyah
University as partial fulfillment the requirement of the degree of
Bachelor of Science in Mathematics

By

Fatima Abed Helal Hussain

Supervised

Dr.Qusay Hatem Akkar

MAY 2017

((أولادكم (الأمم) أنا خلقناه من قبل ولم يسبقنا))

خَيْرُ النَّاسِ الْعِلْمِيُّ الْعَظِيمُ

الاهداء

أهدي ثمرة جهدي المتواضع إلى سيدة تغفوا الجنة تحت قدميها إلى أمي
التي تحتضن أسمى بدعوة بعد صلاتها فحققت ما أصبوا إليه بفضل الله وفضلها . . .
إلى أخوي الذين دعماني من بداية مشواري .
إلى كل من علمني حرفاً وهيئني لأتم دراستي أساتذتي الكرام
وإلى كل من كان له فضلاً بعد الله عليّ .

الشكر والتقدير

لا يسعني وأنا أضع هذه اللمسات الأخيرة في هذه الدراسة

ألا أن أتقدم بالشكر والعرفان لكل أساتذتي الذين تلمذت على أيديهم ولا سيما

أستاذي

الدكتور قصي حاتم عكار

الذي أشرف على بحثي لن تفيه أي كلمات حقه فلولا مثابرته ودعمه المستمر ما تم عملي

ولكل من كانت له مساهمة في طريقي ولرفيقات السكن الجامعي

1-1. ABSTRACT.

The aim of this paper is to prove the stability in the sense of Hyers-Ulam stability of a polynomial equation .more precisely , if x is an approximate solution solution of the equation $X^n + \alpha x + \beta = 0$, then there exists an exact solution of the equation near to x .

1-2. INTRODUCTION

The basic problem of the stability of functional equation asks whether an approximate solution of the Cauchy functional equation $f(x+y)=f(x)+f(y)$ can be approximated by a solution of this equation [12] . In 1940

, S. M. Ulam [16] posed the following problem concerning the stability of functional equations:

Give conditions in order for a linear mapping near an approximately linear mapping to exist .

The problem for the case of approximately additive mappings was solved by D . H . Hyers[3] when G_1 and G_2 are Banach spaces .

since then , the stability problems of functional equations have been extensively investigated by several mathematicians (cf. [2] [1] [11] [13]) .

C. Alsina and R. Ger [1] remarked that the differential equation $y' = y$ has the Hyers-Ulam stability . The result of C . Alsina and R . Ger has been generalized by T, S –E. Takahasi and H . Choda[0] , by T. Miura[8], and also by S-E Takahasi, T . miura and S. Miyajima[11]. Furthermore , the result of Hyers-Ulam stability for first –order linear differential equations has been generalized by T. Miura , S. Miyajima [15], and also by S-M. Jung[5,6] .

recently , G . Wng , M. Zhou and L. Sun[17] discussed the Hyers-Ulam stability of the first-order non homogenous linear differential equation.

Motivated by and connected to the results mentioned above and [7], we consider stability problems for a polynomial equation . in this paper , we will investigate the Hyers-Ulam stability of the following polynomial equation :

$$X^n + \alpha X + \beta = 0 \quad (1-1)$$

Where $x \in [-1, 1]$.

2-1 PRELIMINARIES

The Hyers-Ulam stability if there exists a constant $K > 0$ with the following property : for every $\varepsilon > 0$, $y \in [-1,1]$, if

$$|y^n + \alpha y + \beta| \leq \varepsilon$$

Then there exists some $z \in [-1,1]$ satisfying

$$z^n + \alpha z + \beta = 0$$

Such that $|y - z| < K\varepsilon$.

We call such K a Hyers-Ulam stability constant for equation (1.1)

2-2 MAIN RESULTS

Definition (2-3) :

Contraction mapping : or contraction or contractor , on a metric space (M, d) is a function f from M to itself , with the property that there is some no negative real number $0 \leq K < 1$ such that for all X and y in M .

Definition (2-4) :

Complete metric linear space : is to be complete if every Cauchy sequence in X converges in X .

Now , the main result of this work is given in the following theorem .

Theorem (2-5) :

If $|\alpha| > n$, $|\beta| < |\alpha| - 1$ and $y \in [-1,1]$ satisfies the inequality

$$|y^n + \alpha y + \beta| \leq \varepsilon$$

Then there exists a solution $v \in [-1,1]$ of equation (1-1) such that

$$|y - v| \leq K \varepsilon$$

Where $K > 0$ is a constant .

Proof :

Let $\varepsilon > 0$ and $y \in [-1,1]$ such that

$$|y^n + \alpha y + \beta| \leq \varepsilon$$

We will show that there exists a constant K independent of ε and v such that $|y - v| < K \varepsilon$ for some $v \in [-1,1]$ satisfying $X^n + \alpha X + \beta = 0$.

If we set

$$g(x) = \frac{1}{\alpha} (-\beta - x^n), x \in [-1,1]$$

Then

$$|g(x)| = \left| \frac{1}{\alpha} (-\beta - x^n) \right| \leq 1$$

Let $X = [-1,1]$, $d(x, y) = |x - y|$, then (X, d) is a complete metric space , and g map X into X .

Next , we will show that g is a contraction mapping from X to X .

For any $x, y \in X$, one have .

$$\begin{aligned} D(g(x), g(y)) &= \left| \frac{1}{\alpha} (-\beta - x^n) - \frac{1}{\alpha} (-\beta - y^n) \right| \\ &\leq \frac{1}{|\alpha|} |x^n - y^n| \\ &= \frac{1}{|\alpha|} |x - y| |x^{n-1} + x^{n-2}y + x^{n-3}y^2 + \dots + xy^{n-2} \\ &\quad + y^{n-1}| \end{aligned}$$

Since $|\alpha| > n$, $x, y \in [-1, 1]$, $x \neq y$. we obtain

$$d(g(x), g(y)) \leq \gamma d(x, y)$$

Where $\gamma = \frac{n}{|\alpha|} \in (0, 1)$

Thus g is a contraction mapping from X to X , by S. Banach's

Contraction mapping theorem , there exists unique $v \in X$, such that

$$g(v) = v$$

Hence equation (1.1) has a solution on $[-1, 1]$.

Finally , we show that equation (1.1) has the Hyers- Ulam stability .

Let us introduce the abbreviations $K = \frac{1}{|\alpha|(1-\gamma)}$, *then*

$$\begin{aligned} |y - v| &= |y - g(y) + g(y) - g(v)| \\ &\leq |y - g(y)| + |g(y) - g(v)| \\ &\leq \left| y - \frac{1}{\alpha} (-\beta - y^n) \right| + \gamma |y - v| \\ &= \frac{1}{|\alpha|} |y^n + \alpha y + \beta| + \gamma |y - v| \end{aligned}$$

Thus , we come to the inequalities

$$\begin{aligned} |y - v| &\leq \frac{1}{|\alpha|(1-\gamma)} |y^n + \alpha y + \beta| \\ &\leq K\varepsilon \end{aligned}$$

Which completes the proof .

By applying a similar argument of the proof of theorem 2.5 , it is easy to see the following theorem holds .

Theorem (2.6) :

Let (X, d) be a complete metric linear space ,
T be a contraction mapping from X to X , then $(T - I)x = 0$ has
the Hyers-Ulam stability . That is , for every $\epsilon > 0$, if

$$d(Tx - x, 0) \leq \epsilon$$

Then there exists an unique $z \in X$ satisfying

$$Tz - z = 0$$

$$d(x, z) \leq K\epsilon$$

for some $K > 0$.

proof : since (X, d) is a complete metric linear space and
there exist

$\gamma \in (0, 1)$ such that $d(Tx, Ty) \leq \gamma d(x, y)$ for all $x \neq y, x, y \in X$
, by S. Banach's contraction mapping theorem , there exists
unique $z \in X$, such that $Tz = z$, hence equation $Tv - v = 0$ has a
solution on X .

For every $\epsilon > 0$, if $d(Tx - x, 0) \leq \epsilon$ then

$$\begin{aligned} d(x, z) &= d(x - Tx + Tx - Tz, 0) \\ &= d(Tx - x + Tx - Tz) \\ &\leq d(Tx - x, 0) + d(Tx - Tz, 0) \end{aligned}$$

$$\leq \varepsilon + \gamma d(x, z)$$

Thus , we obtain the inequalities

$$d(x, z) \leq \frac{\varepsilon}{(1 - \gamma)}$$

Which completes the proof .

It is easy to see that by the similar technique , we can discuss the Hyers-Ulam stability of the polynomial equation defined on any finite interval $[a, b]$.

Unfortunately , we could not prove the Hyers-Ulam stability of the polynomial equation defined on an infinite interval .

It is an interesting open problem whether the polynomial equation

$a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0 = 0$ has the Hyers-Ulam stability for the case it has some solution in $[a, b]$.

Acknowledgement. This work was supported by National Natural science Foundation of China (10871213) . The authors would like to thank the referee for a number of valuable suggestions regarding a previous version of this paper .

REFERENCES

- [1] C. Alsina and R. Ger, On some inequalities and stability result related to the exponential function, J.Inequal. Appl. 2 (1998) 373-380.
- [2] P.Enflo and M.S. Moslehian. An interview with Themistocles M. Rassias, Banach J.Math. Anal. 1 (2007) , no. 2, 252-260 .
- [3] D.H. Hyers, On the stability of the linear functional equation, Proc. Nat. Acad. Sci. U.S.A. 27 (1941) 222-224.
- [4] D.H. Hyers, G. Isae and Th.M. Rassias, stability of Functional Equations in several variables, Birkhauser, Basel, 1998 .
- [5] S.-M. Jung, Hyers-Ulam stability of linear differential equations of first order , Apple. Math. Lett. 17 (2004) 1135-1140.
- [6] S.-M. Jung, Hyers-Ulam stability of linear differential equations of first order (II) , Apple. Math. Lett.19 (2006) 854-858 .
- [7] S.-M. Jung, A fixed point approach to the stability of aVolterra integral equation , Fixed point Theory Apple. 2007, Art. ID 57064, 9 pp.
- [8] T. Miura, On the Hyers-Ulam stability of a differentiable map , Sci. Math. Japan 55 (2002) 17-24 .
- [9] T. Miura, S. Miyajima and S.-E. Takahasi, A characterization of Hyers-Ulam stability of first order linear differential operators, J. Math. Ana. Apple. 286 (2003) 136-146.
- [10] T. Miura, S.-E. Takahasi and H.Choda, On the Hyers-Ulam stability of real continuous function valued differentiable map, Tokyo J. Math. 24 (2001) 467-476.
- [11] M.S. Moslehian , Ternary derivations , stability and physical aspects , Acta Appl . Math . 100 (2008).no.2.187-199 .
- [12] Gy. pólya and G. Szegő , Aufgaben und Lehrsätze aus der Analysis , vol. I, Springer, Berlin , 1925 .
- [13] Th. M. Rassias , On the stability of functional equations and a problem of Ulam, Acta Appl. Math. 62 (2000) 23-130 .
- [14] S.-E. Taakahasi , T. Miura and S. Miyajima , On the Hyers-Ulam stability of the Banach space-valued differential equation $y'=\lambda y$, Bull . Korean Math . Soc .39 (2002) 309 – 315 .
- [15] S.-E. Taakahasi , H. Takagi , T. Miura and S. Miyajima , the Hyers-Ulam stability constants of first order linear differential operators , J. Math .Anal. Appl . 296 (2004) 403 – 409.
- [16] S.M. Ulam , A Collection of the Mathematical Problems , Interscience , New York , 1960.
- [17] G.Wang , M. Zhou and L.Sun , Hyers-Ulam stability of linear differential equations of first order Appl. Math. Lett .21 (2008) 1024-1028.