

Journal of Pharmaceutical Sciences and Research

Scopus coverage years: from 2009 to Present

Publisher: PharmaInfo Publications

ISSN: 0975-1459

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science

Visit Scopus Journal Metrics »

CiteScore 2016
0.28

SJR 2016
0.158

SNIP 2016
0.391

<http://www.scimagojr.com/journalsearch.php?q=19700174933&tip=sid&clean=0>

<http://www.jpsr.pharmainfo.in/Contact%20us.php>

Acceptance Letter

22 April 2018

Dear Author(s)

Dr./Mr(s). Nabeel Abed Abdul-Reda / University of Al-Qadisiyah, IRAQ

Dr./Mr(s). Salwa Razzaq Abdul Ameer / University of Al-Qadisiyah, IRAQ

Dr./Mr(s). Raad Saad Jihad / Directorate General of Muthanna Education, IRAQ

Greetings, With reference to your article titled:

**SYNTHESIS AND ANTIMICROBIAL STUDYING OF SOME NEW
FORMAZAN DERIVATIVES FROM (8-CHLOROTHEOPHYLLINE)**

We wish to bring to your kind notice the following

- ✓ we acknowledge the receipt of the above mentioned article
- ✓ the above mentioned article(s) has been sent to the reviewer of expert comments
- ✓ The above mentioned article(s) (**JPSR 100S**) have been accepted and it will be published in the July-August/ 2018 V10/No. (6-7) (Journal of Pharmaceutical Sciences and Research (JPSR) ISSN: 0975-1459)

Synthesis, Identification and Study Biological Activity of Some New Chalcone derivatives from 8-Chlorotheophylline

Nabeel A. Abdul-Reda Salwa R. Abdul-Ameer*

Department of Chemistry, College of Science, University of Al-Qadisiyah, Iraq

*Department of Chemistry, College of Education University of Al-Qadisiyah, Iraq

Email: nabeel.a.alradha@qu.edu.iq

*salwa22.razzaq@gmail.com

Abstract:


New derivatives of 8-Chloro theophylline (8-CTh) were synthesized by react 8-CTh and 4-amino acetophenone to prepared 8-(4-acetylphenylamino)-1,3-dimethyl-1*H*-purine-2,6(3*H*,7*H*)-dione then the result reacts with some substituted benzaldehyde derivatives to prepared different chalcone compound then the products were allowed to react with thiourea and hydrazine to give Pyrimidine and pyrazoline derivatives respectively. The reaction was monitored by thin-layer chromatography (TLC) technique. All new compounds were characterized by melting points, elemental analysis, FT-IR, ¹H, ¹³C and 2D-NMR spectroscopy. Antimicrobial activity of these derivatives was also determined.

Keywords: 8-Chlorotheophylline, chalcone, Pyrazoline, Pyrimidine, antimicrobial.

Introduction

Xanthine a purine base produces in most human body tissues and fluids and in other organisms it is produced by purine metabolism. [1] The commonly used methyl xanthine theophylline, caffeine and theobromine are naturally occurring compounds in various plants. [2] These substances are components of most widely consumed beverages, coffee, tea and

cocoa. Theophylline and its salts are used in the therapy of bronchial asthma and chronic obstructive pulmonary disease (COPD) 8-CTh is a purine derivative which is obtained from theophylline [3].




In therapy, 8-CTh is used in combination with antihistamines in order to enhance their action against travel sickness and other hearing labyrinth diseases [4]. In this work prepared some derivatives from 8-CTh by chalcone reaction. Chalcone can be prepared by condensation reaction between different aldehyde compound and an acetophenone in the presence of opportunity base (aldol conditions) [5]. Recently, the heterocyclic compound of different ring with different heteroatoms are important for biologically active and industrial chemical products [6]. Pyrazoline (Py) and Pyrimidine (Pyr) derivatives are among the most prominent heterocyclic compounds, which have an important role in pharmaceutical and agrochemical industries as well as bioactive products because of their significant and wide spectrum of biological activities [7].

EXPERIMENTAL

All materials were of highest purity and supplied by Merck, Sigma Aldrich and Fluka- company. Melting points were measured on a Buchi melting point apparatus B-545 (Buchi Labortechnik AG, Switzerland). Microanalytical data were obtained with a Vario, Elementar apparatus (Shimadzu, Japan). The IR spectra were recorded on Shimadzu Fourier Transform Infra-red spectrophotometer (Model 270), using KBr discs. NMR spectra were recorded on 600 MHz (^1H) and at 100 MHz and (^{13}C) spectrometers (Bruker, Germany) with TMS as the internal standard and on δ scale in ppm. (TLC) was performed on silica gel for (TLC) and spots were visualized by Iodine vapors. The reagents used were of analytical grade while the solvents were purified before use.

Synthesis of 8-(4-acetylphenylamino)-1,3-dimethyl-1*H*-purine-2,6 (3*H*,7*H*)-dione (S1). [8]



8-CTh (0.21g/0.001 mol) it was dissolve with 4-amino acetophenone


(0.135g /0.001mol) in 15 ml of THF and add 1ml of triethyl amine and then reflux reaction for 24 hours and monitored reaction by TLC using the solution (ethyl acetate: n-Hexane) (4:2) the reaction is neutralized by glacial acetic acid and the precipitate obtained was filtered, washed and recrystallized from ethanol. Yield (79%) as light yellow solid. **m.p** = 176-179 °C. **Rf** = 0.6 , **IR** (KBr, cm⁻¹): N-H (3361), C-H_{arom.}(3130), C-H_{ali.}(2983, 2883), C=O (1716), C=O_{ketone} (1700) C=C_{arom} (1696), C=C (1635) C-N (1373), C=N (1519). **¹H-NMR** (600 MHz, DMSO-*d*₆) δ= 3.18 (s, 3H, N₁-CH₃), 3.53 (s, 3H, N₂-CH₃), 9.72 (s, 1H, N₁₀H), 14.54 (s, br, 1H, N₉H), 2.33 (s, 3H, COMe), 7.57-7.60 (d, 2H, C2'+6' *J*= 8.3 Hz), 7.89-7.91 (d, 2H, C3'+5' *J*= 8.3 Hz) **¹³C-NMR** (100 MHz, DMSO *d*₆): δ= 27.3 (N₁-CH₃), 29.8 (N₃-CH₃), 24.6 (Me), 113.9 (C2'+6'), 115.0 (C5), 127.3 (C4'), 131 (C3'+5'), 143.8 (C1'), 149.4 (C4), 151.4 (C=O₂), 156.5 (C=O₆) , 167.3 (C8), 196.5 (C=O).

General procedure for synthesis chalcone (S2-S4). [9,10]

(3.13 g / 0.01 mol) from S1 with different aromatic aldehydes (0.01 mol). The mixture dissolved in 10 ml of alcohol. Sodium hydroxide solution

3ml (10%) was added slowly and the mixture stirred for 2hr until the entire mixture becomes very cloud then the mixture was poured slowly into 20 ml of water with constant stirring and kept in refrigerator for 24 hours. The reaction was monitored by TLC (ethyl acetate: n-Hexane) (4:2). the reaction is neutralized by glacial acetic acid and, the precipitate obtained was filtered, washed and recrystallized from ethanol


8-(4-((E)-3-(4-hydroxy-3-methoxyphenyl) acryloyl) phenylamino)-1,3-dimethyl-1*H*-purine -2,6 (3*H*,7*H*)-dione (S2).



(3.13 g / 0.01 mol) of S1 with Vanillin (1.52g / 0.01mol) Yield (74%) as Brown solid. **m.p** = 185-188 °C. **Rf** = 0.51 ,**FT-IR** (KBr, v, cm⁻¹): 3336 (N-H), 3113, 2999, 2885 (C-H)_{ali}, 1516 (C=C)_{cha}, 1668 (C=O)_{cha}, 3437 (O-H), 1593, 1471 (C=C)_{ar}. 1267 (C-O-C), 3224 (N₁₀-H)_{cha}. **¹H-NMR** (600 MHz,

DMSO- d_6) δ =3.21 (s, 3H, N₁-CH₃), 3.60 (s, 3H, N₃-CH₃), 3.80 (s, 3H, OMe), 7.64, 7.66 (d, 1H, H_b J =15.9 Hz), 8.0 ,8.03 (d, 1H, H_a J = 15.9 Hz), 9.84 (s, 1H, OH), 12.46 (s. br, 1H, N₉H), 9.64 (s. br, 1H, N₁₀H), 6.54-7.44 (m, 7H_{arom}). ¹³C-NMR (100 MHz, DMSO d_6) δ = 27.6 (N₁-CH₃), 29.7 (N₃-CH₃), 55.4 (OMe), 108.8 (C5), 112.6 (C2"), 114.9 (C2'+C6'+C5"), 121.5 (Ca +C6"), 123.3 (C1"), 124.8 (C4'), 130.4 (C3'+C5'), 144.2 (C1'+Cb), 147.1 (C4), 147.3 (C4" C-OH), 150.8 (C3'(C-OMe) ,151.0 (C=O 2), 153.8 (C=O6), 167.1 (C8), 194.8 (C=O) _{cha}. Anal. calc. for C₂₃H₂₁N₅O₅ (447.45): C 61.74, H 4.73, N 15.65. Found: 61.51, H 4.65, N 15.42.


8-(4-((E)-3-(4-chlorophenyl)acryloyl) phenylamino)-1,3-dimethyl-1H-purine-2,6 (3H,7H)-dione (S3).



(3.13 g / 0.01 mol) from S1 with p-chloro benzaldehyde (1.4g / 0.01mol)

Yield (82%) as Yellow solid. **m.p** = 178-181 °C. **R_f** = 0.55 **FT-IR** (KBr, ν , cm⁻¹): 3460 (N-H), 2995, 2875 (C-H)_{ali}, 1589 (C=C) _{cha}, 1665 (C=O)_{cha}, 1055 (C-O), 1016 (C-Cl)_{cha}, 3219 (N₁₀-H). ¹H NMR (600 MHz, DMSO- d_6) δ = 3.17 (s, 3H, N₁-CH₃), 3.33 (s, 3H, N₃-CH₃) , 7.33-7.68 (m,8H. _{arom}),7.72, 7.76 (d,1H, H_b J =15.6 Hz), 7.99,8.03 (d,1H, H_a J =15.6 Hz), 8.69 (s, br, 1H, NH), 9.99 (s,1H, N₉H) ¹³C-NMR (100 MHz, DMSO d_6) δ = 27.3 (N₁-CH₃), 29.8 (N₃-CH₃), 115.0 (C5), 112.7 (C2'+C6'), 121.3 (Ca), 127.3 (C4'), 129.3 (C3"+C5"), 130.6 (C2"+C6"), 131.1 (C3'+C5'), 134.6 (C4" CCl+C1"), 142.7 (C1'), 144.6 (Cb), 149.4 (C4), 151.4 (C=O 2), 156.5 (C=O 6), 166.9 (C8), 192.6 (C=O _{cha}). Anal. calc. for C₂₂H₁₈ClN₅O₃ (435.87): C 60.62, H 4.16, N 16.07. Found: C 60.40, H 4.04, N 15.79.

8-(4-((E)-3-(4-(dimethylamino)phenyl) acryloyl) phenylamino)-1,3-dimethyl-1H-purine 2,6(3H,7H) dione (S4).



(3.13 g / 0.01 mol) from S1 with p-dimethyl amino benzaldehyde (1.49g / 0.01mol) Yield (78%) as Brown solid.

m.p = 189-191 °C. **R_f** = 0.42 **FT-IR**

(KBr, v, cm⁻¹): 3336 (N-H), 3132, 2939, 2858 (C-H)_{ali}, 3000 (C-H) 1540 (C=C)_{cha}, 1658 (C=O)_{cha}, 1604 (C=C)

¹H-NMR: (600 MHz, DMSO-*d*₆) δ

= 3.38 (s, 3H, N₁-CH₃), 3.21 (s, 3H, N₃-CH₃), 3.06 (s, 3H, NMe₂), 7.50, 7.51 (d, 1H, H_b), 8.06, 8.08 (d, 1H, H_a), 6.54-6.56 (d, 2H, C3''+C5''), 7.60-7.62 (d, 2H, C3'+C5'), 7.64-7.66 (d, 2H, C2'+C6'), 7.72-7.74 (C2''+C6''), 9.84 (s, 1H, N₉H), 12.46 (s, br, 1H, NH).

¹³C-NMR (100 MHz, DMSO-*d*₆) δ = 27.3 (N₁-CH₃), 29.8 (N₃-CH₃), 43.5 (NMe₂), 112.5 (C5), 111.9 (C2'+C6'), 122.9 (C_a), 129.4 (C4'), 111.5 (C3''+C5''), 129.7 (C2''+C6''), 130.1 (C3'+C5'), 151.4 (C4''), 123.3 (C1'') 141.4 (C1'), 143.9 (C_b), 148.1

(C4), 150.1 (C=O 2), 156.2 (C=O 6),

167.2 (C8), 191.5 (C=O)_{cha}.


General procedure for synthesis

Pyrimidine compounds derivatives (S5-S7).[11].

(0.001mol) of prepared chalcone was dissolved in ethanolic sodium hydroxide (10ml) and (0.001mol) was add slowly from thiourea, were stirred about 3-4 hours with a magnetic stirrer.

This was then poured into 5 ml of cold water with continuous stirring for an hour and then kept in refrigerator for 24 hours, the reaction was monitored by TLC (ethyl acetate: n-Hexane) (4:2). The precipitate obtained was filtered, washed and recrystallized with ethanol.


8-(4-(6-(4-hydroxy-3-methoxyphenyl)-3,6-dihydro-2-mercapto pyrimidin-4-yl) phenyl amino)-1,3-dimethyl-1*H*-purine-2,6(3*H*,7*H*)-dione (S5).



(0.447g / 0.001 mol) from S2 with thiourea (0.076 g / 0.001 mol) Yield


(72%) as light Brown solid. **m.p** = 195-199 °C. **R_f** = 0.35, **FT-IR** (KBr, ν , cm^{-1}), 2644 (S-H), 3394 (O-H), 1593 ($\text{C}=\text{C}$)_{ar}, 2993 ($\text{C}-\text{H}$)_{ar}, 1099 ($\text{C}-\text{O}-\text{C}$), 1523 ($\text{C}=\text{N}$)_{pyr}, 1648 ($\text{C}=\text{C}$)_{pyr}. **¹H-NMR** (600 MHz, DMSO-*d*₆) δ = 1.36 (s, 1H, SH) 3.34 (s, 3H, N₁-CH₃), 3.50 (s, 3H, N₃-CH₃), 3.93 (s, 3H, OMe), 4.78 (s, br, 1H, C_{5''}), 5.20 (s, br, 1H, C_{2''}) 6.68 (s, br, 1H, C_{6'''}), 6.88 (s, 1H, br, C_{5''}), 6.96 (s, br, 1H, C_{2''}), 7.56 (s, 1H, C_{3'+5'}), 7.78 (s, br, 1H, C_{2'+6'}) 8.18 (s, 1H, NH_{pyr}), 9.07 (s, br, 1H, N₉H), 9.66 (s, 1H, OH), 11.38 (s, 1H, N₁₀H). **¹³C-NMR** (100 MHz, DMSO *d*₆): δ = 27.6 (N₁-CH₃), 29.7 (N₃-CH₃), 55.4 (OMe), 62.7 (C_{2''}), 103.4 (C_{5''}), 108.1 (C₅), 112.3 (C_{2'''}), 114.9 (C_{5'''}), 121.5 (C_{2'+C6'}), 123.3 (C_{6'''}), 130.4 (C_{3'+C5'}), 125.9 (C_{4'}), 133.4 (C_{1'''}), 136.5 (C_{1'}), 142.3 (C_{6''}), 148.8 (C_{3'''}+C_{4'''}), 149.7 (C₄), 150.8 (C=O 2), 153.6 (C=O 6), 164.6 (C_{4''}), 167.1 (C₈). Anal. calc. for C₂₄H₂₅N₇O₄S (507.57): C 56.79, H 4.96, N 19.32. Found: C 56.56, H 4.82, N 19.09.

8-(4-(6-(4-chlorophenyl)-3,6-dihydro-2-mercaptopyrimidin-4-yl)phenylamino)-1,3-dimethyl-1*H*-purine-2,6(3*H*,7*H*)-dione (S6).



(0.435 g / 0.001 mol) from S3 with thiourea (0.076g / 0.001mol) Yield (69%) as Brown solid. **m.p** = 190-192 °C. **R_f** = 0.29, **FT-IR** (KBr, ν , cm^{-1}): 2653 (S-H), 1604 ($\text{C}=\text{C}$)_{ar}, 1552 ($\text{C}=\text{N}$), 3084 ($\text{C}-\text{H}$)_{ar}, 1523 ($\text{C}=\text{N}$)_{pyr}, 1652 ($\text{C}=\text{C}$)_{pyr}. **¹H-NMR** (600 MHz, DMSO-*d*₆) δ = 1.41 (s, 1H, SH), 3.20 (s, 3H, N₁-CH₃), 3.36 (s, 3H, N₃-CH₃), 4.65 (s, 1H, C_{5''}), 5.23 (s, 1H, C_{2''}), 7.09-7.13 (d, 1H, C_{2'''}+C_{6'''} J = 7.9 Hz), 7.21-7.24 (d, 1H, C_{3'+5'} J = 8.2 Hz), 7.41-7.44 (d, 1H, C_{3'''}+C_{5'''} J = 7.9 Hz), 7.57-7.60 (d, 1H, C_{2'+6'} J = 8.2 Hz), 8.71 (s, 1H, NH), 9.34 (s, br, 1H, N₉H), 10.38 (s, 1H, N₁₀H). **¹³C-NMR** (100 MHz, DMSO *d*₆) δ = 27.5 (N₁-CH₃), 29.7 (N₃-CH₃), 65.4 (C_{2''}), 102.9 (C_{5''}), 110.6 (C₅), 120.8 (C_{2'+C6'}), 123.9 (C_{3'''}+C_{5'''}), 128.7 (C_{3'+C5'}), 129.4 (C_{2'''}+C_{6'''}), 130.1 (C_{4'}), 131.1 (C_{1'''}+C_{4'''}), 135.6 (C_{1'}), 142.9 (C_{6''}), 146.6 (C₄), 151.4 (C=O 2), 153.8 (C=O 6), 159.4 (C_{4''}), 166.8 (C₈) .

8-(4-(6-(4-(dimethyl amino) phenyl)-3,6-dihydro-2-mercaptopyrimidin-4yl)phenylamino)-1,3-dimethyl-1H-purine-2,6(3H,7H)-dione (S7)




(0.444g / 0.001 mol) from S4 with thiourea (0.076 g / 0.001 mol) Yield (78%) as Brown solid. **m.p** = 198-200 °C. **R_f** = 0.25 ,**FT-IR** (KBr, ν , cm^{-1}): 2646 (S-H) 1604 (C=C) ar 3050 (C-H) ar. 1500 (C=N)pyr. , 1649 (C=C)pyr. **¹H-NMR** (600 MHz, DMSO-*d*₆) δ = 1.37 (s, 1H, SH), 3.16 (s, 6H, NMe₂) 3.20 (s, 3H, N1-CH₃), 3.32 (s, 3H, N3-CH₃), 4.68 (s, 1H, C5''), 5.25 (s, 1H, C2'') 6.87-6.89 (d, 2H, C2''' + C6''' *J* = 7.8 Hz), 7.09-7.13 (d, 2H, C3' + 5' *J* = 8.2 Hz), 7.22-7.24 (d, 2H, C3''' + 5''' *J* = 7.8 Hz), 7.34-7.36 (d, 2H, C2' + 6' *J* = 8.2 Hz), 9.57 (s, 1H, NH), 9.76 (s, 1H, N₉H), 12.41 (s, 1H, NH). **¹³C-NMR** (100 MHz, DMSO *d*₆): δ = 27.7 (N1-CH₃), 29.8 (N3-CH₃), 43.6 (NMe₂) , 65.8 (C2''), 103.6 (C5''), 111.1 (C3''' + C5'''), 112.6 (C5),

120.6 (C2' + C6'), 124.1 (C1'''), 128.7 (C3' + C5'), 129.4 (C2''' + C6'''), 130.8 (C4'), 135.6 (C1'), 142.9 (C6''), 144.4 (C4), 146.4 (C4'''), 152.6 (C=O 2), 154.8 (C=O 6), 159.6 (C4''), 167.5 (C8).

General procedure for synthesis pyrazoline compounds derivatives (S8-S10) [12].

A mixture a prepared chalcone (0.001 mol), and (0.001 mol) and hydrazine hydrate (99%) (0.001 mol) in (25 ml) ethanol was refluxed with stirring about 5 hrs, the reaction was monitored by TLC (ethyl acetate: n-Hexane) (4:2). The precipitate obtained was filtered, washed and recrystallized with ethanol.


8-(4-(4,5-dihydro-5-(4-hydroxy-3-methoxyphenyl)-1H-pyrazoline-3-yl)phenylamino)-1,3-dimethyl-1H-purine-2,6(3H,7H)-dione. (S8)



(0.447 g / 0.001 mol) from S2 with hydrazine (0.032g / 0.001mol) Yield (80%) as brown solid. **m.p** = 191-193 °C. **R_f** = 0.38 **FT-IR** (KBr, ν , cm^{-1}): 2960, 2926, 2879 (C-H) _{ali}, 3363, 3219 (N-H) _{Py}, 3039 ,1595 (C=C)_{ar}, 3069

(C-H)_{ar.Py} , 3462 (N-H), 3564 (O-H).
¹H NMR (600 MHz, DMSO-*d*₆) = 3.22 (s, 3H, N₁-CH₃), 3.37 (s, 3H, N₃-CH₃) 3.63 (s, 3H, OMe), 3.78 (s, 1H, C₅ py) , 3.82 (s, 1H, C₄ py), 6.77-7.83 (d, 8H. arom), 9.23 (s ,1H, NH py.), 10.29 (s, 1H, N₉H), 13.38 (s1H, NH), 9.50 (s, 1H, OH), ¹³C-NMR (100 MHz, DMSO *d*₆): δ= 27.6 (N₁-CH₃), 29.7 (N₃-CH₃), 42.4 (C₄Py), 58.8 (OMe), 49.7 (C₅py), 108.8 (C₅), 112.6 (C₂"), 114.9 (C₂' + C₆' + C₅"), 121.5 (C₆"), 123.3 C₄' , 130.4 (C₃' + C₅'), 137.5 (C₁"), 142.6 (C₁'), 147.5 (C₄" C-OH) , 147.7 (C₄), 149.7 (C₃"), 150.8 (C₃py) , 150.0 (C=O 2 + C₃' (C-OMe), 151.0 (C=O 6) , 168.2(C₈).


8-(4-(5-(4-chlorophenyl)-4,5-dihydro-1H-pyrazoline-3-yl)phenylamino)-1,3-dimethyl-1H-purine-2,6(3H,7H)-dione (S9)



(0.435 g / 0.001 mol) from S3 with hydrazine (0.032g / 0.001mol) Yield (77%) as Yellow solid. **m.p** = 189-193 °C. **R_f** = 0.42 **FT-IR** (KBr, ν, cm⁻¹): 3130, 2981, 2926, 2879 (C-H) ali, 3429

(N-H) , 3346, 3227 (N-H) py , 1597 (C=C) ar., 3462 (N-H), 3059 (C-H) ar.
¹H-NMR (600 MHz, DMSO-*d*₆) δ = 3.21 (s, 3H, N₁-CH₃), 3.37 (s, 3H, N₃-CH₃), 3.87 (d, 1H, C₅), 7.43-7.85 (m, 8H. arom), 9.15 (s ,1H, NH py.), 10.09 (s, 1H, N₉H), 3.71 (s, 1H, C₄ py) . ¹³C-NMR (100 MHz, DMSO *d*₆): δ= 27.6 (N₁-CH₃), 29.7 (N₃-CH₃), 108.5 (C₅), 113.1 (C₂' + C₆'), 123.4 (C₄'), 127.5 (C₃' + C₅' + C₂" + C₆"), 129.4 (C₃" + C₅"), 141.4 (C₁"), 134.0 (C₄"), 136.9 (C₁'), 147.2 (C₄), 150.2 (C₃), 48.6 (C₅Py), 151.7 (C₃py) , 42.3 (C₄ py), 150.8 (C=O 2), 153.7 (C=O 6), 167.1 (C₈)

8-(4-(5-(4-(dimethylamino) phenyl)-4,5-dihydro-1H-pyrazoline-3-yl)phenylamino)-1,3-dimethyl-1H-purine -2,6(3H,7H) dione. (S10)



(0.444g / 0.001mol,) from S4 with hydrazine (0.032g / 0.001m) Yield (76%) as brown solid. **m.p** = 199-102 °C. **R_f** = 0.61 **FT-IR** (KBr, ν, cm⁻¹), 3334, 3223 (N-H) py , 1597 (C=C) ar, 1560 (C=N) 3039 (C-H) ar.Py , 3334 (N-H)_{py}, 2953, 2922, 2856 (C-H)_{ali}. ¹H-

NMR (600 MHz, DMSO- d_6) δ = 2.98 (s, 3H, N(Me)₂), 3.20 (s, 3H, N₁-CH₃), 3.36 (s, 3H, N₃-CH₃), 3.47 (s, 1H, C4_{py}), 3.82 (s, 1H, C5_{py}), 6.57-6.59 (d, 2H, C3''+5'' J=8.6 Hz), 6.74-6.77 (d, 2H, C2'+6' J=8.5 Hz), 7.32-7.34 (d, 2H, C2''+6'' J=8.6 Hz), 7.61-7.64 (d, 2H, C3'+5' J=8.5 Hz), 8.36 (s, 1H, NH_{py}), 10.60 (s, 1H, N₉H), 13.06 (s, 1H, NH) 5.98 (s, 1H, C₄_{py}). **¹³C-NMR** (100 MHz, DMSO d_6): δ = 27.5 (N₁-CH₃), 29.7 (N₃-CH₃), 45.5 (NMe)₂, 43.0 (C4_{py}), 49.9 (C5)_{py}, 108.2 (C5), 113.0 (C3''+C5''), 118.1 (C2'+C6'), 125.7 (C4'), 127.5 (C3'+C5'), 127.8 (C2''+C6''), 133.5 (C1''), 138.7 (C1'), 147.3 (C4), 149.0 (C4''), 150.2 (C3)_{py}, 151.4 (C=O₂), 153.4 (C=O₆), 166.8 (C8). Anal. calc. for C₂₄H₂₄N₈O₂ (456.20): 63.15, H 5.30, N 24.55. Found: C 62.92, H 5.21, N 24.31

Antimicrobial Evolution

The newly synthesized compounds were selected for their antimicrobial activities against different bacteria and fungi. The microorganisms used were *Staphylococcus aureus* (Gram positive), *Escherichia coli*, (Gram negative) and *Candida albicans* by using the agar diffusion method [13] to select the most potent compounds. 5 mg of each compound was dissolved in


dimethyl sulfoxide (DMSO, 1 mL) then complete up to 10 mL with distal water to give a concentration of 500 µg/mL. The bacteria were maintained on Muller hinton agar media, the dishes incubated at 37 °C for 24 hr for bacteria while 72 hr for fungal. [14,15] The efficiency of the tested compounds was compared to that of water and DMSO, zone of inhibition measured by ruler.

RESULTS AND DISCUSSION

8-CTh was a starting material for the synthesis of new chalcone compound over a reaction with (P-amino acetophenone) scheme [1]


The appearance of stretching band of secondary aromatic amine (NH) at (3361) cm⁻¹ and disappearance of C-Cl of 8-CTh at (760) cm⁻¹, and showed ¹H-NMR spectrum for N-H 14.54 (s, br, 1H, NH), are attributed to the formation amine compound. ¹³C-NMR spectrum for 24.6 (Me), 113.9 (C2'+6'), 127.3 (C4'), 131 (C3'+5'), 143.8 (C1') [16]. The reaction of S1 compound with different aromatic aldehydes in the presence NaOH to form chalcone in scheme 2 showed appearance of the band 1516, 1589 and 1525 to (C=C)_{cha} for S2, S3 and S4 respectively, in other hand 1668, 1665 and 1658 to (C=O)_{cha} for

S2, S3 and S4 respectively. ^1H -NMR showed the deplete of $\text{CH}=\text{CH}$ proton at about 7.64, 7.66 for Hb, 8.0, 8.03 for Ha to S2 and 7.72, 7.76 Hb, 7.99, 8.03 Ha to S3 and 7.50, 7.51 for Hb, 8.06, 8.08 for Ha to S4. ^{13}C -NMR spectrum for $(\text{C}=\text{O})_{\text{cha}}$. 194.8, 192.6 and 191.5 for S2, S3 and S4 respectively [17]. Some of prepared compounds studding the 2D-NMR (HMBC) and we found for compound S2 three coupling for type $^{1,3}J_{\text{H}, \text{C}}$ the first; between C6" to be the distance $\delta=120$ ppm and proton Hb $\delta=7.5$ ppm, and the second between C1" for the distance $\delta=124$ ppm and proton Ha $\delta=8$ ppm. Third of coupling $^{1,3}J_{\text{H}, \text{C}}$ between C=O for the distance $\delta=195$ ppm and proton Hb for the distance $\delta=7.5$ ppm. or the distance $\delta=195$ ppm and proton Ha for the distance $\delta=8$ ppm.




For S3 also studding the HMBC-NMR and we found three coupling for type $^{1,3}J_{\text{H}, \text{C}}$ the first; between C6" to be the distance $\delta=115$ ppm and proton Hb $\delta=7.5$ ppm, and the second between

C1" for the distance $\delta=135$ ppm and proton Ha $\delta=8$ ppm. Third of coupling $^{1,3}J_{\text{H}, \text{C}}$ between C=O for the distance $\delta=194$ ppm and proton Hb for the distance $\delta=7.5$ ppm. The second type of coupling is $^{1,2}J_{\text{H}, \text{C}}$ there are two coupling between C=O for the distance $\delta=194$ ppm and proton Ha for the distance $\delta=8$ ppm and between C4" for the distance $\delta=135$ ppm and proton H3" for the distance $\delta=7.3$ ppm. [18,19]. The second type of coupling is $^{1,2}J_{\text{H}, \text{C}}$ there are one coupling between C=O for the distance $\delta=195$ ppm and proton Ha for the distance $\delta=8$ ppm.




of Chalcone (S2-S4) that reacted with thiourea in the presence ethanolic Sodium hydroxide to give corresponding. Pyrimidine as well as reacted with hydrazine in the presence absolute ethanol to give pyrazoline derivatives. Pyrimidine and pyrazoline can be prepared by Scheme 3. ^1H -NMR and ^{13}C -NMR of Pyrimidine and

pyrazoline compounds can be show in
Table [1] and [2] respectively.



Scheme 1: Synthesis of 8-(4-acetyl phenyl amino)-1,3-dimethyl-1H-purine-2,6 (3H,7H) –dione.




Scheme 2: Synthesis of chalcone

Table [1]: Characteristic FTIR absorption, ¹H-NMR and ¹³C-NMR bands of compounds (S5-S7)

| No. of comp. | IR | ¹ H-NMR | | | ¹³ C-NMR | | |
|--------------------|-------------------|--------------------|------------------|------------------|---------------------|------------------|------------------|
| | N-H _{py} | N-H _{py} | C4 _{py} | C5 _{py} | C3 _{py} | C4 _{py} | C5 _{py} |
| S8 | 3219,3363 | 9.15 | 6.21 | 3.97 | 151.3 | 42.7 | 50.1 |
| S9 | 3346,3227 | 9.15 | 3.71 | 3.78 | 151.7 | 42.3 | 50.0 |
| S10 | 3334, 3223 | 9.63 | 3.75 | 3.90 | 150.5 | 42.9 | 42.3 |

Table [2]: Characteristic FTIR absorption, ¹H-NMR and ¹³C-NMR bands of compounds (S8-S10)

| No.of comp. | IR | | | ¹ H-NMR | | ¹³ C-NMR | | | |
|----------------|------|-------------|-------------|--------------------|------|---------------------|-------|-------|-------|
| | S-H | C=C pyr. | C=N pyr. | N-H | S-H | C2" | C4" | C5" | C6" |
| S5 | 2644 | 1648 | 1523 | 8.18 | 1.36 | 62.7 | 103.4 | 164.6 | 142.3 |
| S6 | 2653 | 1649 | 1523 | 8.71 | 1.41 | 65.4 | 102.9 | 159.4 | 142.9 |
| S7 | 2646 | 1652 | 1500 | 9.68 | 1.37 | 65.8 | 103.6 | 159.6 | 142.9 |



Scheme 3: Synthesis of Pyrimidine and Pyrazoline derivatives

Biological part

Control of microbial population is necessary to prevent show of disease, infection, decomposition, spoilage and contamination and caused by them. The newly synthesized compounds were screened for their antimicrobial activity invitro against bacteria (*Staphylococcus aureues*, *Escherichia coli*) and fungal (*Candida albicans*) as show in Figure 1. The antimicrobial activity results discovered that most of the tested compounds have moderate to strong activity. The most effective compounds are **S3**, **S5** and **S8** When these compounds were compared with the reference compounds (DMSO, distill water) we found that they have an antimicrobial activity higher or almost equal to them. While *candida albicans* not responsive for all compounds.

CONCLUSION

In this study we are reported synthesis of many chalcone derivatives from 8-CTh. The work included preparation of chalcone compounds and then prepared Pyrimidine when react with thiourea, in other hand react with hydrazine to prepared pyrazoline. These derivatives were found active angst bacterial and not response to fungal. These derivatives confirmed from spectral data analysis; FT-IR, H^1 -

NMR, C^{13} -NMR and 2D-NMR.

ACKNOWLEDGMENT

The authors are thankful prof. Dr. Najim A.AL-Masoudi, Chemistry Department, Konstanz University, Germany, for making and analyzer NMR spectra. Also we are really appreciating the help of Miss Zahra khthair, College of education -Biology, AL-Qadisiya University.




Figure 1: Zone of inhibition of compounds against *Staphylococcus aureus*, *Escherichia coli* and *Candida albican*

| No. of Comp. | Gram-positive | Gram-negative | Fungal |
|--------------|------------------|---------------|-------------------------|
| | <i>S. Aureus</i> | <i>E-coli</i> | <i>Candida ablicans</i> |
| S1 | 12 | 18 | - |
| S2 | 14 | 22 | - |
| S3 | 17 | 23 | - |
| S4 | 14 | 25 | - |

| | | | |
|---------------|----|----|---|
| S5 | 13 | 20 | - |
| S6 | 19 | 23 | - |
| S7 | 15 | 17 | - |
| S8 | 16 | 24 | - |
| S9 | 13 | 16 | - |
| S10 | 15 | 18 | - |
| Control/DMSO | 0 | 0 | 0 |
| Distill water | 0 | 0 | 0 |

Zone of inhibition measured in mm: no activity (0.0), very weak activity (< 7 mm), weak activity (7–10), moderate activity (11–15 mm), strong activity (> 15 mm).

References

- [1] Szentmiklosi, A. J; Cseppento, A; Gesztelyi, R; Zsuga, J; Körtvely, A; Harmati G; Nanasi P.P. *Current Medicinal Chemistry*. **2011** .18, 3695-3706.
- [2] Barnes, P. Theophylline. *Pharmaceuticals*, **2010**, 3, 725-747
- [3] Riksen, N.P.; Smits, P.; Rongen, G.A. The cardiovascular effects of methyl xanthines. *Handb. Exp. Pharmacol.*, **2011**, 200, 413-437.
- [4] Mosselhi, A .M; Abdallah, M. A; Metwally, N.H, El-Desoky I. A; Break, L. M. *ARKAT USA, Inc* .**2009**, 1551-7012, 53 -63.
- [5] Mamedov, I, G; Khrustalev, V.N; Bayramov, M.R; Maharmov, A.M .*INDIAN J.CHEM* .**2017**.56B,192-196.
- [6] Solankee, A; Tailor, R. *Chemistry International* **2016**. 2(4) 189-200.
- [7] Tawfiq, M. T; Zaier A. J. *Journal of College of Basic Education*,**2015**.21.18
- [8] Younis, F.M. *College of Basic Education Researchers Journal*. **2011**.11, 711-721.
- [9] Khazaal, M. S. Tomma ,J. H. *IBN AL- HAITHAM J. FOR PURE & APPL. SCI* .**2011**.24.2.
- [10] Hsieh, C.T. *Bioorganic & Medicinal Chemistry Letters* .**2012**. 22, 3912-3915.

- [11] Elarfi.M. J. *Sci. Revs. Chem. Commun.* **2012** .22, 103-107.
- [12] Voskiene, A; Mickevicius, V; and Mikulskiene, G. *ARKIVOC* .**2007**. xv ,303-314.
- [13] Abdalla, M.; Gomha, S; Abd el-aziz, M; Serag, N. *Turk J Chem.***2016**. 40 .441-453.
- [14] Dabholkar, V.V; Ansari, Y.F. *Indian Journal of chemistry.* **2008**. 47. 1758-1761.
- [15] Joshi, V. D; Kshirsagar, M. D.; Singhal, S. *International Journal of ChemTech Research.***2012**. 4,971-975.
- [16] Abdul-Reda N.A; Abass, A. K and Gebur, N. A. *European Journal of Chemistry* .**2016**. 2 ,206-212.
- [17] Savant, S. S, Patel, S.M, Gamit, E. A, HaJiyani, H. U, Patel, S. A, Patel, P. S and Patel, K.C. *Int J Chem Sci.* **2017** ,15. 2 .139.
- [18] Abdul-Rida, N.A;Mohammed, T.I;Al-Masoudi,N.A;Frotscher,M. *Medical chemistry research.***2017**.22.
- [19] Nakanishi,K; Mill Valley, California: university Science Books.**1990**.136