

Parasitic Infection in Wild Rabbits Oryctolagus Cuniculus

Ikhlas A. Marhoon ^{1*}, Khalid Th. Mattar ¹, Fatima I. Mohammad ²

¹ Department of Biology, College of Education, University of Al-Qadisiyah, IRAQ ² Department of Environment, College of Science, University of Al-Qadisiyah, IRAQ

Received 29 April 2018 • Revised 25 July 2018 • Accepted 12 August 2018

ABSTRACT

The purpose of current study is the Biochemical detection of parasitic infection in wild rabbits Oryctolagus cuniculus and illustrate which one of these parasites can infect humans or economical animals to clarify the important role of rabbits in the transmission of these parasites (as a vector and/or a reservoir host). During the period October 2016 to December 2017, a total of 55 samples of wild rabbits were collected from the local animals market in Al-Diwaniyah province - Iraq, of both sexes with different ages. After dissection and examination of all rabbits to look for ectoparasites infestation and endoparasites infection, then these parasites were prepared according to suitable methods. Results shown that 41 rabbits (74.55%) were infected with eight different types of external and internal parasites. Our results showed three species of external parasites (ectoparasites): the rabbit ear mite Psoroptes cuniculi (25.45%), the poultry red mite Dermanyssns gallinae (30.91%) and nymphs of Linguatula serrata which belong to the tongue worms (21.82%); and also we found five species of internal parasites (endoparasites) which were: the protozoan Eimeria sp.(61.82%), Giardia sp.(16.36%) and Cryptosporidium cuniculus (38.18%), and the larva stage cysticercus pisiformis (54.55%) and finally the nematodes Passalurus ambiguous (32.73%). The results in this biochemical study revealed that rabbits can be infected with common parasites which are: D. gallinae (poultry red mite), nymphs of Linguatula serrata, Eimeria sp. and Giardia sp. these parasites can infected anther hosts beside rabbits. So that our results enhance the important role of rabbits in the spreading of parasites and transmission of infection to other animals as well as humans.

Keywords: parasitic infection, ectoparasite, endoparasite, wild rabbits

INTRODUCTION

Wild rabbits *Oryctolagus cuniculus* are an economically important animals and it was widely used in scientific researches that concerned with pharmacology, immunology, and physiology and it also plays an important role in zoonatic diseases [1]. Rabbits are exposed to various parasitic infections during their lifetime. They act as intermediate or final hosts for these infections [2, 3]. Several biochemical studies have been conducted on ecto and/or endoparasitic infection in wild rabbits around the world. [4] indicated existence of larvea *cysticercus pisiformis* whichelong to the tapeworm *Taenia pisiformis* in the peritoneum cavity of rabbits; [5] reported that rabbits act as intermediate host of Linguatula *serrara* which is a species of external arthropods that settle in the dog's respiratory ducts and that the evolutionary stages of these arthropods settle in the large intestine of rabbits, [6] gave a detailed description of the most important tapeworms that can infect pet rabbits. While [7] reported that rabbits were infected with pinworm *Passalurus ambiguus* at rate (49.65%), also [8] reported 12 different species of rabbit worms, the most important of which are the cestodes: *Cittotaenia ctenoides, Cittotaenia pectinata*, Nematodes: *Nematodirus zembrae, Trichostrongylus retortaeformi, Graphidium strigosum,* and the larva *Cysticercus pisiformis*. On the other hand [9] found wild rabbits infected with two types of protozoa *Cryptosporidium cuniculus* and *Giardia duodenalis* at rate 22.38% and 7.41% respectively.

© 2018 by the authors; licensee Modestum Ltd., UK. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/). Ikhlas.abbas@qu.edu.iq (*Correspondence) Khbio7@gmail.com Fatima.mohammad@qu.edu.iq

Marhoon et al. / Parasitic Infection in Wild Rabbits Oryctolagus Cuniculus

Fable 1. Percentage of infection for the total, ecto and endoparasites which isolated from wild rabbits (Total No. = 55)				
Type of parasite	No. positive	% of positive		
Ectoparasite	23	41.82		
Endoparasite	36	65.45		
Total	41	74.55		

Locally, a few biochemical studies and researches conducted on rabbits' parasitic infection. [10] examined 1500 rabbits to identify the most common diseases, they found that the rate of infection of rabbits with eggs of stomach and intestinal worms was 13.3%, oocysts of Eimeria spp. with an infection rate at 28% and two species of mites which causes scabies: Sarcoptic scabiei and Psoroptes sp.

Other biochemical studies have shown infection of domestic rabbits with several endoparasites with a total infection rate 46.66% which were: two types of protozoa: Eimeria magna and E. steidae, one type of cestoda (larvae of Cysticercus pisiformis) and three types of nematodes were recorded: Passalurus ambiguous, Nematodirus leporis, Trichostrongylus retortaeformis [11]. Another study has shown that two types of external arthropods were isolated from rabbits: Dermanyssus gallinae and Cheyletiella parasitivorax [12]. Recently, the importance of rabbits has increased in our countries because they are depending on their meat as a source of food, and some Europe and Arabian countries have established breeding farms for rabbits such as: the United Kingdom, Germany, French, Aspin, Syria and Egypt.

Because of wild /or domesticated rabbits considerable an intermediate and final hosts for many types of parasites, therefore it becomes a threat to the public health of humans and other animals, so that our biochemical study aims to investigate the types of ecto- and endo-parasites which infected rabbits, demonstrate the percentage of these infections and illustrate which one of these parasites can infect humans or their economical animals.

MATERIAL AND METHODES

Collection of Samples

The current biochemical study included 55 samples of wild rabbits which purchase from the local animals market in Al-Diwaniyah province - Iraq of both sexes with different ages, during from October 2016 to December 2017. The samples were placed in cages that made for breeding of laboratory rabbits until their dissection. Number of species, number of parasites which isolated from each sample were recorded.

Examination and Dissection of Rabbits

The rabbits were anesthetized by using chloroform and then rabbit's hear was examined, especially near ear, noise and upper parts of legs, to investigate infestation of ecto-parasites which preservative in 70% ethanol until preparation [13]. After that, all rabbits were dissected and examined to look for internal parasites. After preparation, by using stains such as iodine stain, Acid fast stain (MZN) to reveal some protozoa, all slides were examined microscopically under X40 and X100 [14].

Statistical Analysis

The results were statistically analyzed by using chi-square at $P \le 0.05$ on using SSPS program.

RESULTS AND DISCUSSION

The results of the present biochemical study showed that 41 specimens out of 55 rabbits were infected with both of external and internal parasites with a total infection rate 74.55%. 23 (41.82%) out of 55 rabbits were infected with ectoparasites, and 36 (65.45%) out of 55 rabbits were infected with endoparasites (Table 1).

The results which appeared in Table 1 showed that the total infection rate in the rabbits under biochemical study was 74.55%, which is a high infection rate. This result gives an important health indicator, because the wild rabbits can be considered a source of continuous and transmission of these parasites to other animals besides of humans.

The results in Table 2 showed that wild rabbits infected with eight different species of parasites which were : three types of ectoparasites: Psoroptes cuniculi (rabbit ear mite) (25.45%), Dermanyssns gallinae (poultry red mite) (30.91%) and nymphs of Linguatula serrata which belong to the tongue worm (21.82%) ; and five types of endoparasites which were: the protozoan Eimeria sp. (61.82%), Giardia sp. (16.36%) and Cryptosporidium cuniculus (38.18%), and the larva stage cysticercus pisiformis (54.55%) and the nematodes Passalurus ambiguous (32.73%).

Group of parasite	Species of parasite	No. of infection	% of infection
	Psoroptes cuniculi	14	25.45
Ectoparasites	Dermanyssns gallinae	17	30.91
	Nymph of Linguatula serrata	12	21.82
	Eimeria sp.	34	61.82
	Giardia sp.	9	16.36
Endoparasites	Cryptosporidium cuniculus	21	38.18
	Cysticercus pisiformis	30	54.55
	Passalurus ambiguous	18	32.73

Table 2. Types of ecto and endoparasites which isolated from wild rabbits



Figure 1. The percentage of parasites which isolation from wild rabbits

The percentage of parasitic infections in wild rabbits (**Figure 1**) appeared that Eimeria sp. (61.82%) and *Cysticercus pisiformis* larvae (54.55%) which belonging to *Taenia pisiformis* had the highest infection rate, while the protozoan *Giardia* sp. had the lowest infection rate (16.36%). The results showed significant differences between the highest and lowest rate of infection at $P \le 0.05$.

The results of the current study were concurred with several studies that recorded infection with these parasites in both domesticated and wild rabbits, such as [15]; [16]; [17] and [18] which found rabbits' infection with [19]. Other studies have shown the high rate of infection with *Cysticercus pisiformis* in wild rabbits [20], as well as many studies recorded infection with pinworm *P. ambiguous* in rabbits [21]. On the other hand, this results consistent with many studies that have documented high rates of infection in rabbits, such as: [22], while others have reported low rates of infection such as [23] which recorded infection between this biochemical study and others may be due to several reasons, including: the areas allocated for raising rabbits and their suitability in health conditions, the degree of contamination of the rabbits' food as well as drinking water. These factors affect in an incidence and prevalence of parasitic infection [24].

Also, some parasites have the same sources of transport such as water and food contaminated with infective stages like mature cyst, oocyst and eggs. This increases the possibility of the incidence of common parasitic infection and demonstrates the absence of antagonism between these parasites, especially intestinal parasites, as in the case of infection with [25]. Some studies have pointed out that the incidence of parasitic infections is more likely to be affected by other species of parasites due to the weakened resistance of the animal's body to other pathogens. Some cases reported that infection with intestinal worms helps in the occurrence of chronic infections in intestinal protozoa in the order lagomorpha [26].

Current biochemical study shown some parasites which isolated from rabbits can infect anther hosts, for example [27] which called poultry red mite, is considerably the most damaging ectoparasite of laying chickens, and his prevalence reach to 94% in Belgium, Germany, Netherlands and Austria [28]. Also in some cases, humans may be infected with the nymph of [28] through ingestion of the nymph and this case named linguatulosis or Halzoon syndrome, human infection with this parasite had been reported in different places in Iran [29]. And with regard to the protozoan *Eimeria* sp. and *Giardia* sp. these protozoa can infect other animals like hens, goats, sheeps and buffaloes [30, 31, and 32].

So that, the presence of common parasitic infections between rabbits and other animals as well as humans enhances the important role of rabbits in the spreading of parasites widely, and transmission of infection among them.

REFERENCES

- 1. Pilar H, Francisca A, Monica F, Felipe B, Manuel S. Effects of restocking rabbits on helminthes fauna of wild rabbits' population in the North West Iberian Peninsula. Acta Parasitol. 2001;46(4):306-312.
- 2. Soulsy EJ. Helminthes, Arthropods and Protozoa of domestic aminals. 8th ed. Bailliere, Tindall, London. 1982:14-169.
- 3. Bowman DD, Lynn RC, Eberhand ML, Gearjis A. Parasitology for veterinarians, 8th ed. W B Saunder Company, Philadelphia, London. 1999:352.
- 4. Lapage G. Veterinary parasitological, Oliver and Boyd, London. 1968:1182.
- 5. Flynn RJ. Parasites of laboratory animals. 1st Ed, The lowa state university press Amen. 1973:493-503.
- 6. Weisborth SH, Flatt RE, Kraus AL. The biology of the laboratory rabbit, Academic Press Inc, London. 1974:317-329.
- 7. Borkovcova M. Endoparasites of some species of domestic animals in Tisnorsko district. Mendel University of Agriculture and Forestry Brno. 1999:129.
- Zhang W, Yujuan SH, Wang R, Liu A, Hong L, Li Y, Cao J. Cryptosporidium cuniculus and Giardia duodenalis in Rabbits: Genetic Diversity and Possible Zoonotic Transmission. J. Clin. Microbiol. 2012;7(2):1-5. https://doi.org/10.1371/journal.pone.0031262
- 9. AL-Khafaji NJ, Rhaymah MSH. Common diseases of rabbits in Mosul Iraq. Iraqi J. Vet. Sci. 1995;8(2):271-277.
- 10. Yagoob G, Hossein HD. Prevalence rate of endoparasites in wild rabbits or east-Azerbaijan province, Iran. Annals Biol. Res. 2011;2(6):31-35.
- 11. El-Hawary NMK. Study on parasitic infection in rabbits. Ms. Vet. Sc., Faculty of Veterinary Medicine, Benha University, Egypt. 2009.
- 12. Coles EH..Veterinary clinical pathology. 4th .ed, Philadelphia, W.B. Saunders Company, London. 1986:386-403.
- 13. Levine ND. Protozoan parasites of domestic animals and of man. Burgess Publ. Co. U.S.A. 1961:255. https://doi.org/10.5962/bhl.title.7000
- 14. Gres V, Voza T, Chabaud A, Landau I. Coccidiosis in wild rabbits Oryctolagus cuniculus. France. J. Parasitol. 2003;10:51-57. https://doi.org/10.1051/parasite/2003101p51
- Xiao L, Fayer R. Molecular characterisation of species and genotypes of Cryptosporidium and Giardia and assessment of zoonotic transmission. Int. J. Parasitol. 2008;38:1239-1255. https://doi.org/10.1016/j.ijpara.2008.03.006
- Shi K, Jian F, Lv C, Ning C, Zhang L. Prevalence, genetic characteristics, and zoonotic potential of Cryptosporidium species causing infections in farm rabbits in China. J. Clin. Microbiol. 2010;48:3263-3266. https://doi.org/10.1128/JCM.00819-10
- Yang D, Yongjun Y, Yue X, Huaming NX, Nong G, Wang X, Peng A, Guangyou Y. Genetic Variation of Taenia Pisiformis Collected from Sichuan, China, Based on the Mitochondrial Cytochrome b gene. Korean J. Parasitol. 2013;51(4):449-452. https://doi.org/10.3347/kjp.2013.51.4.449
- 18. Ashmawy KI, El-Sokkary MY, Abu-Akkada SS, Dewair AW. Incidence of Passalurus ambiguous in domestic rabbits in Behera Province, Alexandria Journal of Veterinary Sciences. 2010;30(1):115-120.
- Li S, Cui P, Fang SF, Lin RQ, Zou FC, Zhu XQ. Sequence variability in four mitochondrial genes among rabbit pinworm (Passalurus ambiguus) isolates from different localities in China. Mitochondrial DNA. 2014. https://doi.org/10.3109/19401736.2013.855898
- Robinson G, Wright S, Elwin K, Hadfield SJ, Katzer F. Re-description of Cryptosporidium cuniculus (Apicomplexa: Cryptosporidiidae): Morphology, biology and phylogeny. Int. J. Parasitol. 2010;40:1539-1548. https://doi.org/10.1016/j.ijpara.2010.05.010
- 21. Nolan MJ, Jex AR, Haydon SR, Stevens MA, Gasser RB. Molecular detection of Cryptosporidium cuniculus in rabbits in Australia. Infect. Genet. 2010;1:1187-1179. https://doi.org/10.1016/j.meegid.2010.07.020
- 22. Sparagano OA, George DR, Harrington DW, Giangaspero A. Significance and control of the poultry red mite Dermanyssus gallinae. Annul Rev. Entomol. 2014;59:447–466. https://doi.org/10.1146/annurev-ento-011613-162101
- 23. George DR, Finn RD, Graham KM, Mul M, Maurer V, Valiente-Moro C, Sparagano OA. Should the poultry red mite Dermanyssus gallinae be of wider concern for veterinary medical science? Parasitol. Vectors. 2015;8:178. https://doi.org/10.1186/s13071-015-0768-7
- 24. Maleky FA. Case report of Linguatula serrata in human throat from Tehran, central Iran. Indian J. Med. Sci. 2001;55(8):439–441.

- 25. Siavashi MR, Assmar M, Vatankhah A. Nasopharyngeal pentastomiasis (Halzoon): report of 3 cases, IJMS. 2002;27(4):192.
- Anaraki MG, Mobedi I, Ariaiepour M, Pourmohammadi Z, Zare-Bidaki MA. Case report of nasopharyngeal linguatuliasis in Tehran, Iran and characterization of the isolated Linguatula serrata. Iran J. Parasitol. 2008;3(1):53–55.
- Peek HW, Landman WJM. Coccidiosis in poultry: anticoccidial products, vaccines and other prevention strategies. Vet. Quart. 2011;31(3):143–161. https://doi.org/10.1080/01652176.2011.605247
- Chartier C, Paraud C. Coccidiosis due to Eimeria in sheep and goats, a review. Small Ruminant Research. 2012;103:84–92. https://doi.org/10.1016/j.smallrumres.2011.10.022
- 29. Koutny H, Joachim A, Tichy A, Baumgartner W. Bovine Eimeria species in Austria. J. Parasitol. Res. 2012;110(5):893-901. https://doi.org/10.1007/s00436-011-2715-7
- 30. Bahrami S, Alborzi AR. Prevalence of subclinical coccidiosis in river buffalo calves of southwest of Iran. 2013;58(4):527-530
- Kommuru DS, Barker T, Desai S, Burke JM, Ramsay A, Mueller-Harvey I, Miller JE, Mosjidis JA, Kamisetti M, Terrill TH. Use of pelleted sericea lespedeza (Lespedeza cuneata) for natural control of coccidia and gastrointestinal nematodes in weaned goats. J. Vet. Parasitol. 2014;204(3-4):191–198. https://doi.org/10.1016/j.vetpar.2014.04.017
- 32. Slutzky GM. Biochemistry of parasites. Naturwissenschaften. 1981;68(10):498-500. https://doi.org/10.1007/BF00365372

http://www.eurasianjournals.com